

Cardiovascular Diseases in Patients with Sclero Dermis and the Development of Risk Factors in Them

Raxmatova Dilbar Baxriddinovna
Bukhara State Medical Institute

Abstract: In patients with systemic scleroderma (TSD), cardiovascular pathology and related complications (heart attack, stroke, sudden coronary death) are among the most common causes of early death in autoimmune rheumatic diseases, despite continuous improvements in diagnosis and treatment. In TSD disease, heart damage occurs in 15-35% of cases. Manifested by heart failure, rhythm disturbances, pain syndrome. In rare cases, mitral heart failure develops including mitral stenosis[19]. Primary heart damage in TSD disease can be accompanied by changes that occur mainly in the myocardium, pericardium, and heart valves. Mortality from cardiovascular complications at TSD is 29% [2].

Base words: systemic scleroderma, hypercholesterolemia, rheumatoid factor, s-reactive protein, arterial hypertension, cardiovascular risk. The results of the worldwide mortality assessment showed that in 2016, worldwide mortality from non-infectious diseases (NICS) was 41 million, accounting for 71% of total mortality. In most cases, death was the result of four NICS: cardiovascular disease (17.9 million), cancer (9 million), chronic respiratory disease (3.8 million), and diabetes mellitus (1.6 million) [6].

In Russia, 1 million is paid annually. about 200,000 people die from cardiovascular diseases when taken at the expense of the population. And all over the world, this figure is 17-18 million. constitutes [18].

Diseases of the cardiovascular system are the most prevalent worldwide and remain the leading cause of disability and death. According to most experts, this problem is expected to continue in this trend for several more decades. According to experts from the World Health Organization, 31% of all deaths are caused by diseases of the cardiovascular system [13,38].

A large number of studies carried out among the population show that patients with rheumatoid arthritis, seronegative arthropathy (psoriatic arthritis, ankylosing spondyloarthritis) and rheumatological diseases such as systemic lupus erythematosus have a higher risk of death from cardiovascular diseases and other complications of internal organ diseases [8].

Changes in the cardiovascular system in rheumatological diseases-early Atherothrombosis, arterial hypertension (AG) and the resulting complications (stroke and myocardial infarction) determine the course and prognosis of these diseases. In most RK, the main cause of death is due to the contribution of cardiovascular pathologies. It has been found that atherosclerosis and AG in the development of Sue are not only associated with the risk factors that lead to them, but also with the immune-inflammatory mechanisms in the disease [20].

In TSD disease, heart damage is considered the most significant of all visceral lesions in terms of importance and occurrence, and is also the main cause of sudden death. The results of the

examination show that heart damage in this disease is 70-90%, which is mainly due to damage to the myocardium and pericardium, as well as cases associated with impaired myocardial excitability and conduction activity are common [15]. In some cases, heart damage in TSD patients develops in a secondary manner after acute sclerodermic renal and pulmonary arterial hypertension. In TSD, vasculopathy is characterized by a progressive restructuring of microcirculation, which leads to the development of various signs of cardiovascular damage. Endothelial dysfunction specific to TSD and hemorrheological disorders in TSD serve as risk factors for the early development of atherosclerosis. Several authors speculate that TSD and atherosclerosis have a common pathogenetic mechanism in vascular damage, and that this process leads to macro and microvascular lesions of different manifestations of MYOCARDINIG in TSD disease [2,3,26,33].

Zhu S.Yu. and research of other hammuallifs (2013) found that immunosuppressor therapy (glucocorticosteroids, d penicillamine, cyclophosphamide, methotrexate, azathioprine, cyclosporine) conducted in such patients in combination with a higher risk of developing acute myocardial infarction in TSD li patients did not reduce the risk of developing myocardial infarction [22]

Recently, it has been found that there is a role of cytokines in the development of cardiovascular diseases in RK. SRO is said to be a link between inflammation in the vessels, coagulation, and thrombosis. SRO has been found to have a direct proaterogenic effect on the vascular wall, stimulating the production of cytokines and adgesion molecules. The purpose of this study is to study myocardial pathology and its relationship to the activity of the process in systemic lupus erythematosus (tqb) and systemic scleroderma [20].

Thus, patients with systemic sclerodermia have been found to be associated with higher mortality rates from cardiovascular disease, higher frequency of cardiovascular risk factors (especially YUIK, hypertriglyceridemia), presence of subclinical atherosclerosis, activity of the Atherosclerosis process, contributing factors to disease activity (increased levels of reactive protein s in the blood, il-2, il-6, presence of anti-topoisomerase-1, anti-centromer antibodies). However, the extent to which risk factors occur, their characteristics in patients with TSD, the presence of subclinical atherosclerosis and its apparent manifestation remain highly inaccurate and sometimes contradictory.

The importance of endothelial dysfunction and various cardiovascular risk factors in the formation of atherosclerosis.

Research in recent years has shown that the leading cause of life expectancy in rheumatological diseases is cardiovascular complications associated with atherosclerotic vascular lesions. According to a large number of studies, patients with rheumatoid arthritis and systemic sclerodermia have a much more frequent form of atherosclerosis until clinical signs appear than general populace [14]. These conditions provide the basis for studying the [7] relationship of cardiovascular and autoimmune pathology. However, in systemic sclerodermia, Vascular Pathology occupies the leading place. And the mechanisms of damage to the cardiovascular system and associated complications remain insufficiently studied [1]. Together with this, analyzes carried out in 2015 showed that this pathology is at a high risk of death from vascular damage [19]. These data represent how important it is to study the clinical features of cardiovascular disease in TSD.

The results of the study show that patients with TSD have a high mortality rate from vascular damage against the background of the development of atherosclerosis, and accounted for 28% of all deaths in TSD (3rd among all deaths) [10].

During the subclinical period of atherosclerosis, a thickening of the sleep artery intima media complex (IMK) was observed when Doppler examination of vessels was performed in patients with TSD, and the presence of atherosclerotic plaque in the sleep artery of such patients was found, and this is significant for early detection of the disease and its prognosis [23].

Nordin A. and others [2013] in a study conducted in Stockholm [Sweden] found that the risk of developing atherosclerosis in the YUIK and peripheral arteries was high. It has also been found that a group of TSD li patients with positive centromere antibodies in the body has the highest risk of signs of subclinical atherosclerosis and cardiovascular complications [35]. In Australia, Ngian G.S. and his co-authors' research in 2012 found that TSD li patients had a high probability of co-occurrence of pulmonary hypertension and YUIK, and that the incidence of cardiovascular risk factors such as obesity, hypercholesterinemia, diabetes mellitus did not differ significantly from patients in the control group [34].

In patients with TSD and subclinical atherosclerosis, it is conspicuous that literature data on intima media thickening in the sleeping artery contradicts each other. Masedo R. and other hammuallifes (2012), however, found intima media thickening in the sleeping artery in patients with TSD, but found no Canday correlation between intima media thickening in the sleeping artery and the severity of the disease [31] recent research results provide information on the presence of subclinical atherosclerosis in patients with TSD [11]. Frerichs M. in a study by others [2014], the presence of atherosclerotic plaque in patients with systemic scleroderma and systemic lupus erythematosus in an unobserved state of sleep artery intima media thickness was found in a sleep artery ultrasound examination and provided information about the presence of subclinical atherosclerosis. [25] Schiopu E. and his co-authors found in their research that patients with TSD had higher levels of atheroscleroetic plaques in the sleep artery compared to those in the control guru. There is an atherosclerotic plaque, patients with TSD have high serum levels of inflammatory proteins (il-2, il-6, s reactive protein, plasminogen - 1 activator inhibitors, endoglin, etc. at the level, and these are involved in the pathogenesis of fibrosis and vasculopathy [30].

In the blood of patients with infectious diseases, enzymes such as superoxididismutase [SOD] and glutathioneductase are detected at high values. An increase in the amount of antibodies to these enzymes provides information about the presence of cardiovascular damage in patients with TSD [5].

Determination of sleep artery intima –media thickness, perceived as a "new" cardiovascular risk factor [28]. Research data shows that an increase in the thickness of the sleep artery intima - media determines the prospect of the occurrence of atherosclerosis-related complications of the cardiovascular system (heart attack, stroke) [27].

Pathological changes in the vascular nucleus begin with the activation of endothelial cells, the expression of adgesion molecules followed by capillary necrosis, and their apaptosis, proliferation of the intima wall, occlusion of the vascular nucleus, and finally organ ischemia [9]. Determination of the thickness of the wall of the sleeping artery and the size of the atherosclerotic plaque in it through the UTT examination in the diagnosis of cardiovascular risk is of great importance in predicting cardiovascular disease [30].

TSD li has been found to increase the risk of the origin of cardiovascular disease in different groups of the population due to the hardening of the arterial wall in patients, precisely due to the deposition of calcium in the coronary vessel wall [29].

Arterial hypertension as a risk factor for cardiovascular disease in recurrent sclerodermia

According to many studies, cardiovascular disease in TSD increases the mortality rate from cardiovascular complications. Including arterial hypertension (AG) is observed much more often when taken compared to the general population. At the same time, one of the factors for the formation of cardiac pathology in the autoimmune process was found to be chronic inflammation, and the increase in signs of inflammation is associated with classic cardiovascular risk factors, in particular hypertension. It should be noted that despite the fact that TSD is one of the autoimmune systemic diseases, the pathogenesis of nosology, its complications and comorbid states have significant differences. In TSD, classic inflammation plays less of a role than rheumatoid arthritis and systemic lupus erythematosus, and in the first place are connective

tissue fibrosis, microvascular injuries and vasospasm . Meanwhile, there are very few studies dedicated to studying the relationship between hypertension and inflammation of the "Scleroderma".

Despite the fact that there are not many works dedicated to assessing risk factors in patients with TSD, and the information obtained does not always have one meaning, cardiovascular diseases in this pathology are more common than in the general population and are detected in young patients. Thus, one of the largest analyzes of 168 scientific studies on traditional and non-traditional risk factors in various autoimmune diseases found that AG is much more common in TSD and that chronic inflammation is associated with the formation of cardiovascular pathology.. Considering that cardiovascular pathology is one of the main causes of death in TSD, the relationship between cardiovascular risk factors associated with the traditional cardiovascular risk factor, namely hypertension vatsd, is extremely difficult. The study showed that patients with arterial hypertension have a longer duration of TSD, as well as a correlation between hypertension and TSD activity, specifically with the prevalence of skin injury and increased SRO. This fact can be considered as the pathogenetic basis of the relationship between two diseases –TSD and AG. The data obtained have scientific and practical potential in understanding the processes of formation of cardiovascular pathology at TSD and the therapeutic possibilities of their correction [16, 17].

Due to the rapid development of atherosclerosis in RK, high risk of complications of infection, especially AG, it is good to identify endothelial activation in them in the early stages of the disease. One of the leading methods for determining AG is daily monitoring of blood pressure. Disruption of microcirculation in TSD and the development of fibrosis processes cause visceral lesions, in particular sclerodermic cardiopathy. Determination of cardiac rhythm regularity is a noninvasive informational method of assessing the state of the mechanisms of neurogumoral regulation of the heart, and is an muxim for determining and preventing the risk of developing infectious complications [21].

References

1. Rakhmatova D. B., Rakhmatova D. B. " Main" Symptoms and leading clinical options for the flow of acute coronary syndromes in women //Asian Journal of Multidimensional Research (AJMR). – 2019. – Т. 8. – №. 11. – С. 69-74.
2. Rakhmatova D. B. Analysis of the risk factors of Chd in persons over 60 years among the population of the city of Bukhara //Asian studies. Индия. – 2019. – Т. 1. – С. 33-38.
3. Rakhmatova D. B., Zikrillaev F. A. DETERMINE THE VALUE OF RISK FACTORS FOR MYOCARDIAL INFARCTION //FAN, TA'LIM, MADANIYAT VA INNOVATSIYA. – 2022. – Т. 1. – №. 4. – С. 23-28.
4. Рахматова Д. Б., Аслонова М. Р. Современные методы медицинской реабилитации пациентов с артериальной гипертензией //Биология и интегративная медицина. – 2018. – №. 3. – С. 110-117.
5. Рахматова Д. Б. и др. ОЦЕНКА ФАКТОРОВ РИСКА ИБС У ЛИЦ СТАРШЕ 60 ЛЕТ СРЕДИ НАСЕЛЕНИЯ ГОРОДА БУХАРЫ //Теория и практика современной науки. – 2018. – №. 5 (35). – С. 704-708.
6. Рахматова Д. Б., Раджабова Г. Х. Анализ факторов риска ИБС у лиц старше 60 лет среди населения города Бухары //Биология и интегративная медицина. – 2019. – №. 3 (31). – С. 37-42.
7. Рахматова Д. Б. Гранат как лечебное средство в народной и древней медицине //Биология и интегративная медицина. – 2022. – №. 1 (54). – С. 157-168.

8. Rakhmatova D. B., Rakhmatova D. B. " Main" Symptoms and leading clinical options for the flow of acute coronary syndromes in women //Asian Journal of Multidimensional Research (AJMR). – 2019. – Т. 8. – №. 11. – С. 69-74.
9. Rakhmatova D. B. Scientific and practical significance of acute myocardial infarction among the population of elderly and old age //Globalization, the State and the Individual. – 2022. – Т. 29. – №. 1. – С. 84-89.
10. Urinov R. M. Therapeutic Possibilities for the Correction of Cognitive and Psychoemotional Impairments in Patients with Post-Covid Syndrome //Tuijin Jishu/Journal of Propulsion Technology. – 2023. – Т. 44. – №. 2.
11. O'rinov R. M., Po'latov S. S. FEATURES OF THE FUNCTIONAL STATE OF AUTONOMIC REGULATION IN PATIENTS WITH POST-COVID SYNDROME //Art of Medicine. International Medical Scientific Journal. – 2023. – Т. 3. – №. 3.
12. Рамазанова Ш. Ш. ГИПЕРМЕТРОПИЧЕСКОЙ АМБЛИОПИИ: АКТУАЛЬНЫЕ ВОПРОСЫ ПАТОГЕНЕЗА И КЛИНИЧЕСКИХ ПРОЯВЛЕНИЙ //BOSHQARUV VA ETIKA QOIDALARI ONLAYN ILMİY JURNALI. – 2023. – Т. 3. – №. 10. – С. 32-38.
13. қизиl Рамазанова Ш. Ш. АМБЛИОПИЯНИ ДАВОЛАШ УСУЛЛАРИНИ ТАКОМИЛЛАШТИРИШ //Евразийский журнал медицинских и естественных наук. – 2022. – Т. 2. – №. 1. – С. 15-16.
14. Каримова С. Ш. Психосоциальная Помощь Больным Расстройством Личности В Период Реабилитации //AMALIY VA TIBBIYOT FANLARI ILMİY JURNALI. – 2023. – Т. 2. – №. 5. – С. 384-386.
15. Каримова С. Ш. КЛИНИЧЕСКАЯ КАРТИНА ЭПИЛЕПТИЧЕСКОЙ БОЛЕЗНИ НА РАЗНЫХ СТАДИЯХ СУДОРОЖНЫХ ПРИСТУПОВ //TA'LIM VA RIVOJLANISH TANLILI ONLAYN ILMİY JURNALI. – 2023. – Т. 3. – №. 5. – С. 313-318.
16. Каримова С. Ш. Психосоциальная Помощь Больным Расстройством Личности В Период Реабилитации //AMALIY VA TIBBIYOT FANLARI ILMİY JURNALI. – 2023. – Т. 2. – №. 5. – С. 384-386.
17. Ruziyev Z. M. COVID-19 SARS-CoV-2 infeksiyasi bilan og'rigan bemorlarda gemostaz tizimining o'zgarish xususiyatlari //Ta'lim fidoylari. – 2022. – Т. 3. – С. 17-23.
18. Kadirovna R. D. Immunological Indicators of Blood and Saliva in Diseases of the Oral Cavity Mucosa in Patients with Covid-19 Complicated by Stage I and II AG //Central Asian Journal of Medical and Natural Science. – 2023. – Т. 4. – №. 2. – С. 43-46.
19. Kadirovna R. D. Indicators of the hemostasis system in the blood of patients with COVID-19 complicated hypertension of both stages //International Conference on Multidimensional Research and Innovative Technological Analyses. – 2022. – С. 179-181.
20. Kadirovna R. D. Immunological Indicators of Blood and Saliva in Diseases of the Oral Cavity Mucosa in Patients with Covid-19 Complicated by Stage I and II AG //Central Asian Journal of Medical and Natural Science. – 2023. – Т. 4. – №. 2. – С. 43-46.
21. Джураев Ш. Д. Факторы Риска Суицидального Поведения При Аффективных Расстройствах У Взрослых И Подростков //Scientific progress. – 2022. – Т. 3. – №. 4. – С. 66-74.
22. Mukhamadeevich R. Z. Features of the Course of Iron Deficiency Anemia in Children //AMALIY VA TIBBIYOT FANLARI ILMİY JURNALI. – 2023. – Т. 2. – №. 5. – С. 266-269.

23. Nabiyeva Z. CLINICAL MANIFESTATIONS OF CHRONIC DISEASES ОРГАНОВ OF THE DIGESTIVE SYSTEM IN CHILDREN //Иновационные исследования в современном мире: теория и практика. – 2023. – Т. 2. – №. 15. – С. 27-28.
24. Sulstonova N. A. THE PROBLEM OF ADDICTED MISSING OF PREGNANCY IN EARLY STAGES OF PREGNANCY //Oriental Journal of Academic and Multidisciplinary Research. – 2023. – Т. 1. – №. 1. – С. 94-101.
25. Sulstonova N. A. Dopplerometric Features of Blood Flow Changes in the Utero-Placental System in Women With Related Pregnancy Mission //Miasto Przyszłości. – 2023. – Т. 34. – С. 268-273.
26. Sulstonova N. A. Evaluation of Clinical and Instrumental Results of Patients with a Risk of Development of Recurrent Mission //Central Asian Journal of Medical and Natural Science. – 2023. – Т. 4. – №. 2. – С. 536-542.
27. Юлдашова Р. У., Наврузова Л. Х. Отношение студентов и преподавателей к электронному обучению //Педагогический профессионализм в образовании. – 2015. – С. 219-220.
28. Urinbaevna Y. R. Features of Prediction of the Severity of Iron Deficiency in Helicobacter Pylori Infection //Scholastic: Journal of Natural and Medical Education. – 2023. – Т. 2. – №. 4. – С. 93-99.
29. Юлдашова Р. У., Жарылкасынова Г. Ж. Анализ Эффективности Препаратов Двухвалентного И Трехвалентного Железа Среди Больных Железодефицитной Анемии В Узбекистане //Central Asian Journal of Medical and Natural Science. – 2021. – С. 437-441.
30. Тиллоева Ш. Ш., Давлатов С. С. Эффективность и переносимость локсидола в лечение ревматоидного артрита у пациентов старших возрастных групп //Central Asian Journal of Medical and Natural Science. – 2021. – С. 432-436.
31. Тиллоева Ш. Ш. и др. Estimation of the condition of the cardiorespiratory system of patients with the concilation of bronchial asthma and arterial hypertension, effects of complex therapy //Новый день в медицине. – 2020. – №. 2. – С. 227-230.
32. Tillaeva S. S. et al. Currency and diagnostic criteria of rheumatoid arthritis in patients of senior age groups //Asian Journal of Multidimensional Research (AJMR). – 2018. – Т. 7. – №. 11. – С. 184-188.
33. Tolibov D. S., Rakhimbaeva G. S. The Seventh European Conference on Biology and Medical Sciences //E31-E33. Austria, Vienna. – 2015.
34. Tolibov, D. S. "Neuropsychological features of Alzheimer's disease." *Vestn. TMA* 2 (2013): 72-76.
35. Tolibov D. S., Rakhimbaeva G. S. The Seventh European Conference on Biology and Medical Sciences //E31-E33. Austria, Vienna. – 2015.
36. Абдуллаева, Ч. А., and У. К. Камилова. "Взаимосвязь процессов ремоделирования сердца с дисфункцией эндотелия у больных с хронической сердечной недостаточностью." *Кардиоваскулярная терапия и профилактика* 15.1 (2016): 16-19.
37. Абдуллаева, Ч. А., and У. К. Камилова. "Взаимосвязь процессов ремоделирования сердца с дисфункцией эндотелия у больных с хронической сердечной недостаточностью." *Кардиоваскулярная терапия и профилактика* 15.1 (2016): 16-19.
38. Jurayev A. M. RJ Khalimov New methods for surgical Treatment of Perthes Disease in children International Journal of Psychosocial Rehabilitation, Vol 24, Issue 02, 2020.

39. DJuraev A. M., Khalimov R. J. New methods for surgical treatment of perthes disease in children //International Journal of Psychosocial Rehabilitation. – 2020. – Т. 24. – №. 2. – С. 301-307.
40. Рахматуллаев Х. Р., Джураев А. М., Халимов Р. Д. Хирургическое лечение болезни Пертеса у детей //В сборнике статей" Турнеровские чтения. – 2020. – Т. 54. – С. 304-307.
41. Джураев Ш. Д. Факторы Риска Суицидального Поведения При Аффективных Расстройствах У Взрослых И Подростков //Scientific progress. – 2022. – Т. 3. – №. 4. – С. 66-74.
42. Жўраев Ш. Ж. Методы Лечения, Основанные На Клинических Особенности Суицида При Эндогенно-Аффективных Расстройствах //Barqarorlik Va Yetakchi Tadqiqotlar Onlayn Ilmiy Jurnalı. – 2022. – Т. 2. – №. 11. – С. 98-101.
43. Жураев Ш. Ж. Специфические Особенности Развития И Течения Шизофрении У Больных Женского Пола //Ta'lim Va Rivojlanish Tahlili Onlayn Ilmiy Jurnalı. – 2022. – Т. 2. – №. 12. – С. 322-326.
44. Jumanazarovich J. S. ENDOGEN AFFEKTIV BUZILISHDA SUITSIDNING KLINIK XUSUSIYATLARI ASOSIDA DAVOLASH CHORA-TADBIRLARI //BOSHQARUV VA ETIKA QOIDALARI ONLAYN ILMİY JURNALI. – 2023. – Т. 3. – №. 6. – С. 1-6.
45. Жураев Ш. Ж. Клиническая, Психосоматическая Характеристика Назогенного Нервно-Психопатологического Состояния У Больных Некоторыми Онкологическими Заболеваниями //Central Asian Journal of Medical and Natural Science. – 2023. – Т. 4. – №. 6. – С. 463-466.
46. Афакова М. СОВРЕМЕННЫЕ ПРЕДСТАВЛЕНИЯ ЭТИО-ПАТОГЕНЕЗА РАЗВИТИЯ КАРИЕСА ПОСТОЯННЫХ ЗУБОВ У ДЕТЕЙ ШКОЛЬНОГО ВОЗРАСТА //International Bulletin of Medical Sciences and Clinical Research. – 2023. – Т. 3. – №. 6. – С. 29-34.
47. Муртазаев С., Афакова М. СРОКИ ПРОРЕЗЫВАНИЯ И МИНЕРАЛИЗАЦИИ ПОСТОЯННЫХ ЗУБОВ У ДЕТЕЙ ШКОЛЬНОГО ВОЗРАСТА //Stomatologiya. – 2020. – Т. 1. – №. 2 (79). – С. 83-88.
48. Джаббарова М. Б. Распространенность и клинические проявления бронхиальной астмы //Биология и интегративная медицина. – 2021. – №. 1 (48). – С. 160-171.
49. Раджабова Г. Б., Джаббарова М. Б., Саломова Н. К. Меры по профилактике факторов риска хронической обструктивной болезни легких //Новый день в медицине. – 2020. – №. 4. – С. 519-521.
50. Джаббарова М. Б. и др. Сравнительная характеристика лекарственных средств, используемых для лечения хронических гепатитов и цирроза печени //Новый день в медицине. – 2019. – №. 4. – С. 151-154.
51. Джаббарова М. Б. и др. Особенности профилактики артериальной гипертензии у подростков //Врач-аспирант. – 2007. – Т. 16. – №. 1. – С. 54-56.
52. Мустафаева Ш. А. Функционально-Морфологическое Состояние Клеток Почек У Больных Ревматоидным Артритом //AMALIY VA TIBBIYOT FANLARI ILMİY JURNALI. – 2022. – Т. 1. – №. 4. – С. 22-27.
53. Мустафаева Ш. А. Функционально-Морфологическое Состояние Почек При Ревматоидном Артритом //AMALIY VA TIBBIYOT FANLARI ILMİY JURNALI. – 2022. – Т. 1. – №. 3. – С. 19-24.

54. Мустафаева Ш. А. Применение лечебной физической культуры в лечении функциональных состояний нервной системы у спортсменов (обзорная статья) //Вестник Совета молодых учёных и специалистов Челябинской области. – 2016. – Т. 5. – №. 4 (15). – С. 34-38.
55. Рахматова М. Р., Жалолова В. З. The place of innovative technologies in training of highly qualified personnel in the highest medical educational institutions //Биология и интегративная медицина. – 2018. – №. 3. – С. 234-247.
56. Rylova N. V. Actual aspects of studying athlete's body composition //Kazan medical journal. – 2014. – Т. 95. – №. 1. – С. 108-111.
57. Рахматова М. Р., Жалолова В. З. Юниор ва кадет спортсменларда тананинг композицион таркибини ўрганиш //Тиббиётда янги кун. – Т. 2. – №. 30/2. – С. 67-70.
58. Рахматова М. Р., Жалолова В. З. Effectiveness of the combined application of interactive methods" debats" and" a weak link" in the conduct of the lesson //Биология и интегративная медицина. – 2018. – №. 4. – С. 225-231.
59. РАХМАТОВА М. Р., ЖАЛОЛОВА В. З. EFFECTIVENESS OF THE COMBINED APPLICATION OF INTERACTIVE METHODS" DEBATS" AND" A WEAK LINK" IN THE CONDUCT OF THE LESSON //Биология и интегративная медицина. – 2018. – №. 4. – С. 225-231.
60. Rakhmatova M. R. et al. The level of knowledge of students acquired in interactive ways" Blitz method" and" Case study". – 2019.
61. Жалолова В. З. и др. Роль инновационных методов обучения на развитие уровня знаний студентов //Новый день в медицине. – 2019. – Т. 4. – №. 28. – С. 32-35.
62. ЖАЛОЛОВА В. З., РАХМАТОВА М. Р. Anthropometric indicators of juniors and cadets in sport medicine //Биология и интегративная медицина. – 2020. – №. 4. – С. 5-15.
63. Zamirovna J. V. Methods for Selecting Junior and Cadets Athletes by Morphofunctional Criteria //Central Asian Journal of Medical and Natural Science. – 2021. – С. 87-91.
64. Рахматова М. Р., Жалолова В. З. Effectiveness of the combined application of interactive methods" debats" and" a weak link" in the conduct of the lesson //Биология и интегративная медицина. – 2018. – №. 4. – С. 225-231.
65. DJuraev A. M., Khalimov R. J. New methods for surgical treatment of perthes disease in children //International Journal of Psychosocial Rehabilitation. – 2020. – Т. 24. – №. 2. – С. 301-307.
66. DJuraev A. M., Khalimov R. J. New methods for surgical treatment of perthes disease in children //International Journal of Psychosocial Rehabilitation. – 2020. – Т. 24. – №. 2. – С. 301-307.
67. DJuraev A. M., Khalimov R. J. New methods for surgical treatment of perthes disease in children //International Journal of Psychosocial Rehabilitation. – 2020. – Т. 24. – №. 2. – С. 301-307.
68. Джураев А. и др. Наш опыт хирургического лечения врожденного возвышения лопатки у детей раннего возраста //Медицина и инновации. – 2021. – Т. 1. – №. 4. – С. 37-44.
69. Мухамедова Ш. Т. и др. Внутрибольничная инфекция у новорожденных детей //Биология и интегративная медицина. – 2021. – №. 3 (50). – С. 75-86.
70. Мухамедова Ш. Т. Особенности динамики цитокинов у новорожденных с синдромом системного воспалительного ответа. – 2020.

71. Давронова Х. З. Снижение коэффициента фракционного анизотропии как фактор развития когнитивных нарушений у постинсультных больных //Actual Issues and Solution of Development of Economic Sectors of the Republic of Uzbekistan in Modern Conditions Proceedings of the international conference.–Jizzakh. – 2022. – Т. 1. – С. 764-770.
72. Ходжиева Г. С. Интразональность и специфика течения функциональных заболеваний билиарного тракта при синдроме Жильбера //Научный форум: Медицина, биология и химия. – 2018. – С. 64-68.
73. Ходжиева Г. С. Интразональность и специфика течения функциональных заболеваний билиарного тракта при синдроме Жильбера //Научный форум: Медицина, биология и химия. – 2018. – С. 64-68.
74. Орзиев З. М., Ходжиева Г. С. Диапазон факторов экстрагепатических" субтрансаминаземий" //Биология и интегративная медицина. – 2018. – №. 4. – С. 50-61.
75. Ходжиева Г. С. ЗНАЧЕНИЕ ОБРАЗОВАТЕЛЬНО-ПЕДАГОГИЧЕСКИХ ТЕХНОЛОГИЙ В ФОРМИРОВАНИИ КЛИНИЧЕСКИХ ЗНАНИЙ УЧАЩИХСЯ //BARQARORLIK VA YETAKSHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – Т. 2. – №. 12. – С. 793-798.