

Energy Drinks May Affect the Ovarium

Ismatova M. M., Sobirova SH. S.

Bukhara state Medical institute

Abstract: Energy drinks have an impact on concentration levels, physical performance, speed of reaction, and focus, but these drinks cause many adverse effects and intoxication symptoms. The main goal of this study was to determine the effect of energy drink consumption on ovarian reserve and serum anti-mullerian hormone (AMH) levels. Female Wistar albino rats (n=16) were included and randomized into two groups (n=8). Serum AMH levels were checked before and after energy drinks were given. Eight weeks later, the ovaries and uteruses of the rats were analyzed histopathologically. The number of follicles in the ovaries was counted. The total number of the preantral plus small antral follicles, which show the ovarian reserve, was decreased at the end of eight weeks in both the control group and the energy drink group.

Keywords: Energy drinks, anti-mullerian hormone, ovarian reserve, antral follicles.

Introduction. Energy drinks (EDs) have become much more popular since the 1960s. These drinks are categorized as sugar-sweetened beverages. They also contain caffeine, taurine, glucuronolactone, and other vitamins and mineral additives.(1) EDs have been the fastest growing area of the beverage industry to date. In advertisements, companies assert that these drinks have a good impact on concentration levels, physical performance, speed of reaction, focus, and wellness. Despite the positive effects, these drinks cause many cardiovascular adverse effects and intoxication symptoms causing concerns about the health of the consumers. From 2001 to 2008, the level of ED consumption in adolescents and adults was estimated to have increased from 24% to 56%, causing greater concern.(2)EDs have been completely prohibited or sold in low caffeine forms in some countries because of their adverse effects. Turkey is one of the countries that prohibit the high caffeine forms, but in most countries, EDs are qualified as nutrient support and there are no restrictions. The United States of America Food and Drug Administration updated the classification of EDs as dietary supplements.

Animal Maintenance and Treatment. In this study, sixteen healthy adult female albino rats (8 to 10 weeks old) weighing 190 ± 10 g were used. The animals were kept according to the institutional review board's guidelines for animal care, in a 14-hour light cycle at controlled temperatures (22-28 °C), and food and water were available ad libitum. The water consumption of the rats was not recorded. The weight of the rats was recorded daily and the food they consumed was recorded weekly. After the acclimation period, the stages of the estrus cycles of the rats were evaluated by performing daily vaginal smears.(7)The rats were randomly assigned to two study groups (8 rats each). In the control group (group I), the rats were kept on a normal diet and given water for 8 weeks. In the ED group (group II), the rats were kept on a normal diet and given water plus a daily single dose of ED. The dosing was calculated in comparison with the surface area of humans and rats (3.9 mL/kg b.w.)⁴. A 250-mL can of commercially available ED (A-5330 Fuschl am See, Austria) was opened daily between 09:00 and 10:00 and approximately 0.7 mL was given orally for each rat via flexible oral gavage tubes. A single dose of the ED is roughly equivalent to the minimal human dose [1 can (250 mL)/day], but of course

it varies according to the animal's surface area. Each 100 mL of ED contains a mixture of water, taurine (0.4%) (400 mg), caffeine (0.032%) (320 mg/L), gluconolactone (0.24%), inositol, sucrose, glucose, sodium citrate, carbon dioxide niacin (8 mg), pantothenic acid (2 mg), vitamin B6 (2 mg), B12 (0.002 mg), caramel, riboflavin, and natural and artificial flavoring and coloring agents (these are listed ingredients on the label).

Blood Sampling, Tissue Collection, and Histopathologic Analysis After the acclimation at the beginning of the study (initial) and following the 8-week period (final), blood samples (1 mL) were obtained from the right jugular vein of each rat to measure the serum AMH levels under general anesthesia. The animals were anesthetized by administering 50 mg/kg 10% ketamine hydrochloride (Ketazol; Richter Pharma) and 5 mg/kg 2% xylazine (Rompun; Bayer Healthcare) intramuscularly. All blood samples were immediately centrifuged at 4000 g for 10 minutes, and the collected sera were transferred to Eppendorf tubes. The samples were then transferred on ice and kept at -80 °C in a deep freeze until analysis using an automatic enzyme-linked immunosorbent assay (ELISA) system with a commercially available kit (Cusabio Biotech Co, Wuhan, China). The AMH assay measured concentrations with an assay range of 0.2-15 ng/mL; the manufacturer-specific mean inter and intra-assay coefficient of variation (CV) was less than 15% (CV<15%). All samples and standards were assayed in duplicate as recommended in the catalogue of AMH. The rats were sacrificed in estrus cycles using cervical dislocation and bilateral oophorectomies and hysterectomies were performed in all rats. Histologic ovarian and uterus tissue samples were evaluated by a single histopathologist who was blinded to the origin of the samples. The volumes of the ovaries and uteruses were measured under a microscope. Tissues were fixed in 10% formaldehyde for 72 hours, underwent routine tissue processing, and then embedded in paraffin wax. Four-micron-thick sections were taken from the tissues and the tissues were completely consumed. All sections were stained with hematoxylin and eosin. All sections were investigated under a light microscope (Zeiss Axioskop 40 Carl Zeiss Göttingen, Germany) and the pieces were photographed (AxioVision 3.1 Zeiss Axioplan 2 imaging Germany, Göttingen). Descriptive statistics for the studied variables (characteristics) are presented as median, mean, standard deviation, minimum, and maximum values.(17,18) The Mann-Whitney U test was performed to compare the groups. A statistical significance level was considered as 5% and the Statistical Package for the Social Sciences (SPSS) (Ver. 22) statistical program was used for all statistical computations. There was statistical significance between the means of the weight changes of the two groups (p=0.002). The weekly consumed pellets of the rats showed a statistical difference between group I and group II in regards to food consumption at the end of the 8 weeks .The uterus (n=16) and ovarian tissues (n=32) excised from the rats after fourteen cycles were evaluated morphologically.(19.) When the ovarian tissues of the rats that were given EDs were compared with the control group, their mean ovarian volume was smaller ($10.78 \pm 2.9 \text{ mm}^3$) but there was no statistically significant difference (p=0.99). The endometrium, myometrium, and serosa layers of the uteruses of both groups were histologically normal. (20)Endometrial volumes (mm^3) of both groups were evaluated in stereology in terms of endometrium thickness. The mean volume of endometrium in the ED group (38.83 ± 21.2) was more than in the control group (29.28 ± 14.48) According to these findings, the ED that was used in the experimental group did not affect the endometrium or other layers of the uterus. The total follicles in the ovaries were evaluated. There was no statistically significant difference between the follicles of the two groups (p=0.283). In terms of ovarian reserve analysis, the number of PF, SF, and PA plus small antral follicles (PSF) were counted. Even though the means of PF and SF were decreased more in the subject group, there was no statistically significant difference between the two groups (p=0.026, p=0.057). Furthermore, the means of the total number of PSFs were decreased more in the subject group (160 ± 30.9) than in the control group (133 ± 28.6) and a statistical significance was shown between the two groups.

Discussion. Ovarian function is very important for reproductive health. Follicles play a key role in the reproductive function in the development of follicles, along with several local factors, systemic (hypothalamus and/or pituitary) mediators affect their functions at a certain stage.(34)

AMH is the most important mediator indicating the functions of follicles and providing information about their reserves. PF and AF counts are responsible for the synthesis of this mediator. It was reported that AMH levels were a better indicator for ovarian reserve than age, follicle-stimulating hormone (FSH), luteinizing hormone (LH), inhibin-B, and, estradiol (E₂). AMH is a dimeric glycoprotein that belongs to the transforming growth factor family. AMH protein expression starts immediately after the follicle recruitment and continues to the antral stage of the follicle. PFs are the main source of folliculogenesis in the ovaries. As soon as PF development begins, AMH plays a protective role by slowing down the rate of consumption of the local primordial follicle pool from the granulosa cells. AMH also regulates the growth rate of follicles by inhibiting FSH-related follicle growth in the early antral period. Three-quarters of all AMH is found in PFs and SA follicles. In follicles without AMH in rats, preliminary estrus cycle loss was observed due to the rapid depletion of the primordial pool. Loss of PFs causes irreversible infertility. The common active ingredients used in EDs are caffeine, taurine, sugar, and a vitamin complex. The biggest difference between these drinks and sugary drinks is that they contain caffeine and taurine.(32,33) Taurine is an important amino acid that has many functions in the body and is found in many .Taurine is found in oocytes, granulosa, and theca cells, especially in the epithelial cells of the ovary and uterus. It was reported that cells achieve this through cysteine sulfinic acid. However, there is mRNA-carrying taurine in the ovarie. It was reported that in rats, taurine stimulated follicular development indirectly by the release of FSH, LH, and E₂ through the hypothalamic-pituitary axis, or directly through E₂ produced in granulosa cells by increasing androgen synthesis in the osmoregulator or theca .In an *in vitro* study, it was reported that taurine could directly stimulate follicular development, as well as work as an osmotic regulator in embryos, mouse and human oocytes, and maintain the development of follicles and embryo. In another *in vitro* study on rats, it was suggested that taurine directly stimulated the development of follicles through several ambiguous way.(28,29) In an *in vitro* study on cattle, it was concluded that taurine was not useful in the embryonal development of bovine oocytes as a direct effect. In this study, we propose that the hormonal effect and growth affect the PSF follicle pool, which is the highest, the number of follicles due to stimulation in the PSF pool decreases (p=0.021) and accelerates the preovulatory oocyte passage with a decrease in AMH. We suggest that PFs, whose functions are inhibited, do not provide a reduced PSF pool despite the decrease in AMH, and taurine does not affect PFs because it has no intracellular function. In our study, the number of large AFs increased in subjects given ED. However, we suggest that taurine accelerated the development of oocytes that entered the growth cycle and increased the number of preovulatory oocytes due to the increased LF count (group I; 16±9.4, group II; 17.1±10.7), even though there was no statistical difference (p=0.283). This study is an animal experiment and there is no literature on the dose for ED use. Another difficulty of the study is that there are many components in EDs. The limitation of our study in the histologic examination is that immunohistochemical staining showing PF activity could not be performed. The effect of EDs on fertility could not be evaluated because the reproduction of the rats during the study period could not be controlled.

References

1. Muxiddinovna, I. M. (2022). IMPACT OF ENERGY DRINKS AND THEIR COMBINATION WITH ALCOHOL TO THE RATS METOBOLISM. *Gospodarka i Innowacje.*, 22, 544-549.
2. Mukhiddinovna, I. M. (2022). EFFECTS OF CHRONIC CONSUMPTION OF ENERGY DRINKS ON LIVER AND KIDNEY OF EXPERIMENTAL RATS. *International Journal of Philosophical Studies and Social Sciences*, 2(4), 6-11.
3. Muxiddinovna, I. M. (2022). Effects of Energy Drinks on Biochemical and Sperm Parameters in Albino Rats. *CENTRAL ASIAN JOURNAL OF MEDICAL AND NATURAL SCIENCES*, 3(3), 126-131

4. Ильясов, А. С., & Исмадова, М. М. (2022). ЖИНСИЙ АЪЗОЛАРГА ЭНЕРГЕТИК ИЧИМЛИКЛАРНИНГ САЛБИЙ ТАСИРИ. *Uzbek Scholar Journal*, 5, 66-69.
5. Muxiddinova, I. M. (2022). Damage of Energy Drinks on the Spermatogenesis of Male Rat's. *Research Journal of Trauma and Disability Studies*, 1(9), 111-118.
6. Muxiddinova, I. M. (2022). Effects of Energy Drinks on Biochemical and Sperm Parameters in Albino Rats. *CENTRAL ASIAN JOURNAL OF MEDICAL AND NATURAL SCIENCES*, 3(3), 126-131.
7. Ихтиярова, Г. А., & Розикова, Д. К. (2023). МИКРОБИОЛОГИЧЕСКИЕ ИЗМЕНЕНИЯ У БЕРЕМЕННЫХ С РЕПРОДУКТИВНЫМИ ПОТЕРЯМИ В АНАМНЕЗЕ. Finland International Scientific Journal of Education, Social Science & Humanities, 11(4), 1002-1008.
8. Salimova, T. B. (2022). Features of the Course of Pregnancy in Pregnant Women with Fetal Growth Restriction Syndrome and the Role of Doppler Velocimetry. *Central Asian Journal of Medical and Natural Science*, 3(6), 557-563
9. Salimova, T. B. Homila O'sishi Chegaralanishi Sindromi Bilan Homilador Ayollarda-Homiladorlikning Kechishi Va Uning Diagnostikasida Dopplerometriyaning O'rni.
10. Muxiddinova, I. M. (2022). Ameliorative Effect of Omega-3 on Energy Drinks-Induced Pancreatic Toxicity in Adult Male Albino Rats. *INTERNATIONAL JOURNAL OF HEALTH SYSTEMS AND MEDICAL SCIENCES*, 1(5), 13-18.
11. G. F., M. ., & F. Sh., X. . (2022). Pathomorphological Changes Occurring in the Spleen as a Result of External and Internal Factors. *International journal of health systems and medical sciences*, 1(5), 132–137.
12. Fazliddinova, M. G. . (2023). Oncopsychology of Patients with Breast Cancer after Treatment. *Scholastic: Journal of Natural and Medical Education*, 2(2), 111–116.
13. Razokov, I. B., Rahimov, D. A., Ismatova, M. M., & Kurbonzoda, S. N. MAIN FACTORS CAUSING IRON-DEFICIENT ANEMIA IN CHILDREN OF EARLY AGES. In *SPECIAL EDITION FOR 1st BUKHARA INTERNATIONAL MEDICAL STUDENTS CONFERENCE* (p. 61).
14. Muxiddinova, I. M., & Sobirova, A. Z. (2022). Pregnancy with Twins with Preeclampsia. *Central Asian Journal of Literature, Philosophy and Culture*, 3(11), 212-221.
15. U., N. A., & F., M. G. (2023). Morphofunctional Changes of the Spleen Under the Influence of Various Factors in Postnatal Ontogenesis. *Web of Semantic: Universal Journal on Innovative Education*, 2(5), 228–233.
16. F., M. G., & U., N. A. (2023). Age-Related Morphological and Morphometric Parameters of the Spleen in Postnatal Ontogenesis. *Web of Semantic: Universal Journal on Innovative Education*, 2(5), 234–242
17. Muxiddinova, I. M., & Sobirova, A. Z. (2022). Anemia Iron Deficiency in Pregnancy. *Central Asian Journal of Literature, Philosophy and Culture*, 3(11), 191-199.
18. Mukhiddinova, I. M. (2022). ENERGY DRINKS MAY AFFECT THE OVARIAN RESERVE AND SERUM ANTI-MULLERIAN HORMONE LEVELS IN A RAT MODEL. *BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI*, 2(12), 626-632.
19. Makhmudova Guljamol Fazliddinova. (2023). Age- related factors in diagnosis of malignant neoplasms of the female genital area. *IQRO JURNALI*, 2(2), 675–682.

20. Нурбобоев Адхамбек Уйгунович, Махмудова Гулжамол Фазлиддиновна. (2023). Возрастные особенности в диагностики опухолей молочной железы. *IQRO JURNALI*, 2(2), 683–688.
21. Mukhiddinovna, I. M. (2023). High Caffeine Exposure Increases Ovarian Estradiol Production in Immature Rats. *JOURNAL OF HEALTHCARE AND LIFE-SCIENCE RESEARCH*, 2(3), 8-11.