

Isoniazid-Resistant Mycobacterium Tuberculosis: Frequency of Detection, Resistance Spectra and Genetic Determinants of Resistance

Tosheva Dilnoza Rakhmatovna

Assistant, Department of Microbiology, Virology and Immunology, Bukhara State Medical Institute, Bukhara, Republic of Uzbekistan

Abstract: The lack of rapid diagnosis of tuberculosis with pathogen resistance to isoniazid with preserved sensitivity to rifampicin (IR-TB) may be the reason for low effectiveness of therapy and lead to amplification of drug resistance, including the formation of multidrug resistance. The aim of the work was to determine the incidence of IR-TB in the modern population, to characterize the phenotypic sensitivity and genetic determinants of resistance to isoniazid in representatives of this group of *M. tuberculosis* using representative material. We analyzed the results of drug susceptibility testing obtained from the study of *M. tuberculosis* isolates / *M. tuberculosis* DNA isolated from patients with pulmonary tuberculosis from the clinical departments of the Central Research Institute of Tuberculosis using molecular genetic and/or cultural methods for the period 2020–2023. The incidence of IR-TB was 10% of all detected cases of tuberculosis. *M. tuberculosis* with IR were both monoresistant to isoniazid (40%) and multiresistant (resistant to 3–5 antituberculosis drugs), and resistance to isoniazid was caused by mutations in the *katG* gene, leading to a high level of resistance. Based on an analysis of literature data and our own observations, the importance of the development and implementation of new simple molecular tests to determine resistance to both rifampicin and isoniazid is emphasized.

Key words: Mycobacterium tuberculosis, isoniazid, *katG* mutations, ethionamide, prothionamide, GeneXpert, MDR, XDR, rifampicin

COMPLIANCE WITH ETHICAL STANDARDS: a retrospective analysis of the results obtained during routine laboratory tests for patients undergoing treatment at the Central Research Institute of Tuberculosis was carried out; All patients signed voluntary informed consent for the study. Drug-resistant tuberculosis (TB) is a major public health problem. The main focus is currently on combating multidrug-resistant tuberculosis (MDR-TB), i.e. tuberculosis in a form in which the pathogen is simultaneously resistant to the two most effective anti-tuberculosis drugs (ATDs) - isoniazid and rifampicin. Russia ranks third in the world in terms of MDR-TB prevalence.

In 2022, the incidence and prevalence rates of MDR-TB in the Russian Federation stabilized and amounted to 1.1 and 21.4 per 100,000 people, respectively. At the same time, the proportion of patients with MDR-TB among bacterial isolates increased (from 24.2% in 2020 to 20.3% in 2021 among new TB cases and from 50.0% in 2020 to 55.3% in 2021 among all patients with respiratory TB secreting *Mycobacterium tuberculosis*). Against this background, insufficient attention is paid to other forms of resistance. One of these forms, identified by WHO as a

separate group, is isoniazid-resistant tuberculosis (Hr-TB, IR-TB), which is characterized by pathogen resistance to isoniazid and sensitivity to rifampicin.

Isoniazid is a 1st-line drug that is highly effective for the treatment of active tuberculosis and has a bactericidal effect on *M. tuberculosis*. Phenotypic resistance to isoniazid is associated with mutations in a number of genes (*katG*, *inhA*, *ahpC*, etc.), the products of which are involved in the pharmacokinetics and pharmacodynamics of isoniazid in the bacterial cell.

Inadequate therapy for IR-TB creates a high risk of developing acquired drug resistance to other anti-TB drugs, including rifampicin, leading to the development of MDR. According to WHO, the prevalence of IR-TB is 5–11% depending on the region. There is insufficient information on the prevalence of IR-TB in Russia.

Purpose of the study: to characterize the frequency of occurrence of IR *M. tuberculosis* isolated from patients with pulmonary tuberculosis from the clinical departments of the Federal State Budgetary Institution "Central Scientific Research Institute of Infectious Diseases" for the period 2021–2023, to provide an expanded characteristic of phenotypic sensitivity and to describe the genetic determinants of resistance to isoniazid of this group of *M. tuberculosis*.

MATERIALS AND METHODS: We studied clinical isolates of *M. tuberculosis* and/or *M. tuberculosis* DNA isolated from diagnostic material obtained from each patient admitted to the consulting and clinical departments of the Federal State Budgetary Institution "Central Research Institute of IT" (for the period 2020–2023). All types of microbiological studies were carried out from one portion of diagnostic material.

Cultural diagnostics: Detection of *M. tuberculosis* was carried out using Middlebrook 7H9 liquid medium in the BACTEC MGIT 960 system (BD; USA) according to the manufacturer's standard protocol. Phenotypic drug sensitivity was determined by a modified method of proportions in the BACTEC MGIT 960 system (BD; USA) to eight anti-tuberculosis drugs at critical concentrations: isoniazid (H, 0.1 µg/ml), rifampicin (R, 1.0 µg/ml), ethambutol (E, 5.0 µg/ml), pyrazinamide (Z, 100.0 µg/ml), ethionamide (Eto, 5.0 µg/ml), amikacin (Am, 1.0 µg/ml), capreomycin (Cm 2.5 µg/ml) and levofloxacin (Lfx 1.0 µg/ml) according to standard procedures.

RESEARCH RESULTS: Cultural and molecular genetic studies of diagnostic material from 4056 patients with pulmonary tuberculosis were carried out. In 71 cases, neither *M. tuberculosis* DNA nor *M. tuberculosis* culture was isolated; these cases were excluded from the study. From the remaining 3985 samples of diagnostic material, DNA and/or cultures were isolated *M. tuberculosis* and determined phenotypic/genotypic sensitivity to anti-TB drugs. When obtaining results for clinical isolates of *M. tuberculosis* by both cultural and molecular genetic methods (MGM), in case of their discrepancy, priority was given to data obtained by culture. 38 strains for which, during the culture study, resistance to both isoniazid and rifampicin were determined, but mutations in *rpoB*, which determine resistance to rifampicin, were not determined, were classified as MDR, since the tests we used for the molecular genetic determination of DR detect only a limited number of mutations, so genetic determinants of rifampicin resistance may not have been identified in these cases. Conversely, 29 strains with identified mutations in *rpoB*, which did not have the phenotypic manifestation of resistance to rifampicin, were classified as IR. Since cultural methods for diagnosing tuberculosis are less sensitive than molecular genetic methods, some samples did not give culture growth *M. tuberculosis* on nutrient media. Therefore, phenotypic sensitivity to TTPs was determined for only 250 *M. tuberculosis* isolates with IR. When classifying *M. tuberculosis* isolates with IR according to the nature of drug resistance, we relied on the following definitions: monoresistance - resistance of *Mycobacterium tuberculosis* to only one of the anti-TB drugs; multiresistance - resistance of *Mycobacterium tuberculosis* to two or more anti-tuberculosis drugs, but not to a combination of isoniazid and rifampicin.

The frequency of occurrence of resistance to isoniazid with preserved sensitivity to rifampicin was studied in *M. tuberculosis* isolated from patients with pulmonary tuberculosis from the clinical departments of the Federal State Budgetary Institution "TsNIIT" for the period 2021–2023.

The frequency of occurrence of this form of TB and the dynamics of population spread in different regions of the world have their own specifics. Thus, when analyzing data on the drug sensitivity of the tuberculosis pathogen submitted to WHO for the period 1994–2009. From 131 TB facilities around the world, the highest incidence of IR-TB was reported in Eastern Europe (15%), lower in Western and Central Europe (11%), and in other WHO regions did not exceed 8%.

For a number of regions, these authors showed a trend towards a decrease in the prevalence of this form of tuberculosis, for others, on the contrary, an increase, but in most regions no clear linear dynamics of the indicator was established. The frequency of occurrence of IR described by us *M. tuberculosis* (11%) is similar to the indicators characteristic of Eastern Europe, and the dynamics of the indicator, as in most regions, was non-linear. In a systematic review on the relationship between primary resistance to isoniazid and the acquisition of resistance to other anti-drug medications, it was concluded that acquired resistance to any anti-drug (not only transition to MDR) in isoniazid-mono-resistant strains occurs 5.1 times more often compared to drug-resistant strains. - sensitive strains. The high frequency of occurrence of multidrug-resistant strains shown in our study (55% of all IR), resistant except for isoniazid in addition to 1–5 anti-TB drugs, confirms the possibility of amplification of drug resistance in *M. tuberculosis* with IR.

In this regard, two clinical studies should be mentioned, the purpose of which was to establish the dependence of the effectiveness of therapy with high doses of isoniazid on IR forms of TB on mutations in the genome of the pathogen. It is known that mutations in *katG* (dominated in our study) lead to a high level of resistance to isoniazid, and mutations only in *inhA* lead to a low level of resistance. According to the results of clinical studies, isoniazid therapy was effective if the pathogen had mutations at the *inhA* level, and unfavorable treatment outcomes occurred with mutations at the *katG* level.

All of the above indicates that it is necessary to develop effective treatment regimens for IR-TB. In addition, it is important to pay attention to the rapid diagnosis of isoniazid DR. Rapid, specific and sensitive diagnosis of *M. tuberculosis* lymph nodes can only be provided by molecular genetic methods (1–2 days compared to several weeks of culture diagnostics), which also make it possible to obtain information about the mutation and the level of resistance to isoniazid of the pathogen. This makes MGM the most popular in diagnosing tuberculosis and determining the drug resistance of the pathogen.

CONCLUSIONS: Tuberculosis with IR pathogen can be considered as a potential precursor to MDR tuberculosis. It is therefore important to control the spread of primary isoniazid resistance and prevent amplification of resistance. Analysis of the frequency of occurrence of resistance to isoniazid with preserved sensitivity to rifampicin in *M. tuberculosis* isolated from patients with pulmonary tuberculosis showed a fairly high level of occurrence of IR-TB (more than 12% of all analyzed cases) - usually with mutations leading to a high level of resistance to isoniazid. The data obtained emphasize the importance of accelerated determination of the drug sensitivity of *M. tuberculosis* to both rifampicin and isoniazid using molecular genetic methods in laboratories at all levels. To ensure this possibility, it is necessary to develop new simple point-of-care tests that do not place high demands on laboratory infrastructure.

СПИСОК ЛИТЕРАТУРЫ / REFERENCES

1. Rakhmatovna T. D. Influence of Clinical and Immunological Parameters on the Frequency and Prevalence of *Helicobacter Pylori* in Patients with Gastritis //EUROPEAN JOURNAL OF INNOVATION IN NONFORMAL EDUCATION. – 2022. – T. 2. – №. 12. – С. 115-117.

2. Rakhmatovna T. D. Influence of Clinical and Immunological Parameters on the Frequency and Prevalence of Helicobacter Pylori in Patients with Gastritis //EUROPEAN JOURNAL OF INNOVATION IN NONFORMAL EDUCATION. – 2022. – Т. 2. – №. 12. – С. 115-117.
3. Rakhmatovna T. D. Influence of Clinical and Immunological Parameters on the Frequency and Prevalence of Helicobacter Pylori in Patients with Gastritis //EUROPEAN JOURNAL OF INNOVATION IN NONFORMAL EDUCATION. – 2022. – Т. 2. – №. 12. – С. 115-117.
4. Rakhmatovna T. D. Influence of Clinical and Immunological Parameters on the Frequency and Prevalence of Helicobacter Pylori in Patients with Gastritis //EUROPEAN JOURNAL OF INNOVATION IN NONFORMAL EDUCATION. – 2022. – Т. 2. – №. 12. – С. 115-117.
5. Ашуров О. Ш. Эпидемиологические аспекты коронавирусной инфекции и научные рекомендации по лечению коронавирусной инфекции //НАУЧНОИССЛЕДОВАТЕЛЬСКИЙ ЖУРНАЛ ПО ТРАВМЕ И ИНВАЛИДНОСТИ. – 2021. – Т. 1. – №. 5. – С. 37-43.
6. Shavkatovich, Ashurov Otabek. "Medical and Social Aspects of Epidemiological Analysis and Prevention of Tuberculosis Incidence in the Population." *AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI* (2022): 146-148.
7. Rakhmatovna T. D. Influence of Clinical and Immunological Parameters on the Frequency and Prevalence of Helicobacter Pylori in Patients with Gastritis //EUROPEAN JOURNAL OF INNOVATION IN NONFORMAL EDUCATION. – 2022. – Т. 2. – №. 12. – С. 115-117.
8. Ашуров О. Ш. Медико-Социальные Аспекты Эпидемиологического Анализа И Профилактики Заболевания Населения Туберкулезом Легких //AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI. – 2023. – Т. 2. – №. 2. – С. 137-140.
9. Tosheva, Dilnoza. "Causal Relationship of Immuno-Microbiological Parameters in Tuberculosis With Multiple and Broad Drug Resistance." *INTERNATIONAL JOURNAL OF HEALTH SYSTEMS AND MEDICAL SCIENCES* 2.5 (2023): 259-262.
10. Rakhmatovna T. D. Features of Drug Resistance of the Causative Agent of Tuberculosis in Modern Conditions //AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI. – 2023. – Т. 2. – №. 4. – С. 59-63.
11. Тошева Д. Р. Особенности Лекарственной Устойчивости Возбудителя Туберкулеза В Современных Условиях //AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI. – 2023. – Т. 2. – №. 2. – С. 52-58.
12. Rakhmatovna T. D. Features of Clinical Manifestations of Gastroduodenitis Associated With H. Pylori in Children //BOSHQARUV VA ETIKA QOIDALARI ONLAYN ILMIY JURNALI. – 2021. – Т. 1. – №. 6. – С. 112-114.
13. Shavkatovich, Ashurov Otabek. "Characteristics of the Coronavirus Disease and Its Epidemiological Features." *BOSHQARUV VA ETIKA QOIDALARI ONLAYN ILMIY JURNALI* 1.6 (2021): 99-105.
14. Ашуров О. “Бирламчи Ўпка Туберкулёзини Дори Дармонларга Чидамлилигини Аниклашинг Клиник, Эпидемиологик Ва Микробиологик” Жихатлари.
15. Ашуров О. Ш. Медико-Социальные Аспекты Эпидемиологического Анализа И Профилактики Заболевания Населения Туберкулезом Легких //AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI. – 2023. – Т. 2. – №. 2. – С. 137-140.

16. Al-Hajoj S, Varghese B, Shoukri MM, Al-Omari R, Al-Herbwai M, Alrabiah F, Alrajhi AA, Abuljadayel N, Al-Thawadi S, Zumla A, Zignol M, Raviglione MC, Memish Z. Epidemiology of antituberculosis drug resistance in Saudi Arabia: findings of the first national survey. *Antimicrob Agents Chemother.* 2013;57:2161–2166. [PMC free article] [PubMed] [Google Scholar]
17. Araújo AK, Menezes EA, Santos CD, Cunha FA, Campelo CL. Avaliação da resistência de drogas antituberculosas no estado do Ceará no período de 2000-2002. *Rev Bras Anal Clin.* 2005;37:15–18. [Google Scholar]
18. Baptista IMFD, Oelemann MC, Opromolla DVA, Suffys PN. Drug resistance and genotypes of strains of *Mycobacterium tuberculosis* isolated from human immunodeficiency virus-infected and non-infected tuberculosis patients in Bauru, São Paulo, Brazil. *Mem Inst Oswaldo Cruz.* 2002;97:1147–1152. [PubMed] [Google Scholar]
19. Brito RC, Gounder C, Lima DB, Siqueira H, Cavalcanti HR, Pereira MM, Kritski AL. Resistência aos medicamentos anti-tuberculose de cepas de *Mycobacterium tuberculosis* isoladas de pacientes atendidos em hospital geral de referência para tratamento de AIDS no Rio de Janeiro. *J Bras Pneumol.* 2004;30:425–432. [Google Scholar]