

Research of Health Conditions and Genetic Variants of Young Athletes Involved in Mixed Sports (Review Article)

Jalolova V. Z.

Bukhara State Medical Institute

Abstract: In our country, everything is paid to the development of professional sports. The success of athletes in high-level competitions is an important component of the country's international prestige. In this regard, the situation regarding the importance of the health of athletes for their achievement of high sports results has acquired special relevance [4; 17; 19]. Sports genetics is still at the beginning of the path, but at the same time it opens up many prospects for the development of medical and biological support for sports. The use of the genetic characteristics of the organism will lead humanity to new records, because now not only persistence, regular training, willpower and motivation of an athlete are important, but also his "Olympic" heredity. A detailed study of these genes is necessary for the correct organization of the training process, for predicting the capabilities of athletes. According to modern concepts of functional genomics, it is believed that individual differences in the degree of development of certain physical and mental qualities, as well as in the susceptibility to certain diseases of athletes, are largely due to DNA polymorphisms (variable regions in the DNA sequence). The FABP2 gene produces a protein that binds fatty acids in the digestive system, and thus directly regulates the absorption of fats in the epithelial cells of the small intestine (Baier, Sacchettini et al. 1995). Studies have shown that in the presence of the risk allele, the binding of fatty acids from food occurs to a greater extent and, the presence of the risk allele is directly associated with an increased body mass index, body fat, abdominal adipose tissue size and overweight (Baier, Sacchettini et al. 1995).

Keywords: health conditions, sports genetics, predicting the capabilities of athletes, selection of athletes, DNA polymorphism.

One of the main theoretical and applied medico-biological problems of physical culture and sports medicine is the problem of sports selection, the development of the theory of which determines the level of sports achievements and sports science in general. The results of scientific research have proved that the optimal health indicators of athletes are based on the state of dynamic balance between the functional reserves of the body and the factors affecting it [1; 9; 22; 23]. At the same time, the magnitude of the influence of the factors that are inherent in the modern system of training athletes on the body of those involved increases in proportion to the stages of their professional development, which requires not just optimal health indicators, but the presence of an appropriate level of reserves of functions of its components [7; 12; 26]. The value of the latter is the basis of the athlete's reliability - an indicator that is characterized by high performance of actions and its stability in extreme conditions of activity [5]. In this regard, there is a consensus among specialists that athletes-children and adolescents have the lowest reliability coefficient, since it is this contingent of those involved in sports that has certain prerequisites for the development of maladaptive disorders in somatic systems, the structural

elements of which are included in the newly formed system of adaptation of the body to muscular activity [thirteen; 16; 20; 26].

These prerequisites include specific features of the structure of the child's body, biochemical processes and functions in general, as well as individual organs, in which qualitative (development) and quantitative (growth) changes occur at various stages of ontogenesis [2; 12; 18]. In addition, the inconsistency of regulatory systems and the heterochronism of growth and maturation processes only exacerbate the problem of the development and progression of pathological deviations in the somatic systems of the body of young athletes with an irrational structure of the training process, which is confirmed by the data of scientific research on the morbidity of children-athletes [3; eleven; 21]. Regardless of the preventive and therapeutic measures taken, the number of somatic diseases among reserve athletes does not statistically decrease [13; 20; 26], which allows us to state the presence of a number of issues, the solution of which is associated with the development of the concept of physical rehabilitation of young athletes with dysfunctional disorders of the somatic systems, mediated by training and competitive activities.

Scientific substantiation and development of conceptual, organizational and methodological components of this concept is impossible without a detailed study of the health status of athletes of a particular specialization, without taking into account the characteristics of the biological, psychosocial and professional stages of ontogenesis. In the classical practice of sports medicine, it is generally accepted that all preventive, therapeutic and rehabilitation measures should be carried out on the basis of data from an in-depth medical examination of athletes, the resulting component of which is a conclusion on the admission of the examined to training activities [5; 15; 24]. The admission is carried out according to the following criteria: "healthy", "practically healthy", "requires limitation (full or partial) of training activity". But it follows from practical experience that the data obtained are sufficient for carrying out therapeutic and rehabilitation measures for a specific athlete, but not enough to substantiate the organizational and methodological foundations of the system of measures aimed at protecting the health of a vast contingent of athletes, especially during critical periods of ontogenesis.

At present, most of the sports games can be defined as disciplines "mixed energy system" [1], in which effort has intermittent characteristics with the participation of both aerobic and anaerobic metabolic pathways [2]. In this regard, physical fitness is one of the most important elements that affect the performance of athletes from mixed sports due to the high physical load due to intermittent activity [3]. Indeed, strength and endurance can be considered the key fitness elements for mixed sports such as football, wrestling and motorcycling [4, 6]. All of these sports have been classified as mixed in terms of their metabolic predominance, thus showing the general multifaceted nature of physical demands, which include aerobic capacity, strength, endurance, strength and agility [4, 6]. However, descriptive studies indicate that the physiological characteristics of an athlete can vary greatly across these disciplines, suggesting that there are different sport-related profiles in terms of the relationship between endurance and speed. Athletes from mixed sports disciplines do not have a clear focus on endurance or strength, thus demonstrating wide phenotypic heterogeneity [7]. It is common knowledge that genetic factors influence sprint / strength and endurance [21, 23]. , but only a few studies have examined the relationship between genetic factors and athletic performance in mixed athletes. Among the potential polymorphisms potentially associated with muscle function and physical performance, the most studied are angiotensin-converting enzyme (ACE), variant R577X of the actinin-3 gene (ACTN3), and muscle creatine kinase isoform (CK-). MM), peroxisome proliferator-activated receptor α (PPAR α), and more recently SLC2A4 [9, 14]. It has been demonstrated that alternative polymorphisms of these potential polymorphisms are associated with sprint / strength or endurance disciplines with reasonable replication in different groups of athletes [15,18]. Indeed, the I allele of the insertion (I) / deletion (D) polymorphism in the ACE gene is one of the most putative factors determining aerobic capacity in performance-oriented endurance [8]. Likewise, the PPAR-G, ACTN3 X, and CK-MM A alleles are involved in improving aerobic

performance [16, 17, 19]. On the other hand, it is assumed that ACE D, PPARG C, ACTN3 R and CK-MM G are more related to strength-oriented characteristics [8,16,20]. Given that the contribution of specific genes to athletic performance has been investigated primarily in athletes who are at the two endpoints of the human athletic performance continuum, less is known about the application of sports genomics in more complex disciplines that do not exhibit specific phenotypes. Thus, the aim of the present study was to compare the allele and genotype frequencies of four known polymorphisms in athletes from various disciplines of mixed sports in order to investigate the genetic markers suitable for distinguishing between the predominant components of these sports [25,26].

One of the rapidly developing areas of modern genetics is the development of molecular genetic approaches to determine a person's predisposition to various types of activity. So, in particular, in recent years, a search has been carried out for molecular genetic markers that determine a person's ability to perform high sports loads (Montgomery, 2000; Rogozkin, 2004), which is determined by the need to justify a system for selecting people for sports and correcting the training process. This approach is the most promising, since it allows you to determine the genetic predisposition to high physical activity and to carry out a targeted differentiated selection of children for sports at the earliest stages of their sports activity. It should be noted that in 2000 a human genetic map was created, in which genes were introduced, which in at least one study revealed associations with physical indicators and / or influenced human health (Rankinen, Bray et al, 2006). In an early 2000 version, the map included 29 genes. The 2005 version, 6th - amended, includes 165 autosomal genes, 5 - located on the X chromosome, as well as 17 mitochondrial genes. To date, work of this kind is being carried out only in five countries: the USA, Great Britain, Australia, Russia (Scientific Research Institute of Physical Culture - under the leadership of Doctor of Biological Sciences, Professor V.A. Rogozkin; laboratory of molecular genetic research of the Department of Genetics at the Bashkir State Pedagogical University - under the leadership of Doctor of Biological Sciences, Professor Gorbunova V.Yu.) and Kazakhstan. On the site www.genoterra.ru - the register of leading scientific institutions, the laboratory of molecular genetic studies of the Department of Genetics of BSPU is included in the list of groups engaged in the search for genetic markers that determine the predisposition of people to sports activities [27,28].

Thus, it becomes clear that, from the point of view of health savings, it is of particular interest to study not just the health state of young athletes, but a comprehensive level-by-level screening of the functional state of the athletes. This type of layer-by-layer "screening" will allow to determine the corrected risk factors for the development of somatic pathology in athletes within the framework of uncorrected factors (determinants), in this case it is sports specialization and a stage in the process of many years of training.

According to the research of Svetlana Lyugailo (Results of studying the health status of young athletes: the first level of screening studies) _ using classical approaches to assessing the health status of young athletes, on the basis of the existing criteria for their admission to sports, the previously determined trends in the influence of ontogenetic risk factors for the emergence and development of somatic pathology in the age aspect and in the aspect of professional development.

The above organizational miscalculations in the development of the structural components of the concept of physical rehabilitation for dysfunctional disorders of the somatic systems of the body of young athletes (somatic diseases) will be taken into account in the group of exogenous corrected risk factors for diseases in athletes.

Bibliography:

1. Abete, Itziar, et al. "Nutrigenetics and nutrigenomics of caloric restriction." *Progress in molecular biology and translational science* 108 (2011): 323-346.

2. Vimalleswaran, Karani S., et al. "Candidate genes for obesity-susceptibility show enriched association within a large genome-wide association study for BMI." *Human molecular genetics* (2012): dds283.
3. Vanden, Heuvel JP. "Nutrigenomics and nutrigenetics of ω 3 polyunsaturated fatty acids." *Progress in molecular biology and translational science* 108 (2011): 75-112.
4. Fenech, Michael, et al. "Nutrigenetics and nutrigenomics: viewpoints on the current status and applications in nutrition research and practice." *Journal of nutrigenetics and nutrigenomics* 4.2 (2011): 69-89.
5. Weggemans, R. M., et al. "Genetic polymorphisms and lipid response to dietary changes in humans." *European journal of clinical investigation* 31.11 (2001): 950-957.
6. Djordjevic, Natasa, et al. "Induction of CYP1A2 by heavy coffee consumption is associated with the CYP1A2-163C> A polymorphism." *European journal of clinical pharmacology* 66.7 (2010): 697-703.
7. Rasulovna R. M. Method for Assessing Body Composition and Neurophysiological Characteristics of Junior Athletes and Cadets, Taking into Account the Polymorphism of Genes Responsible for Metabolism // *Central Asian Journal of Medical and Natural Science*. – 2021. – С. 131-136.
8. Rakhmatova M.R., Jalolova V.Z., Methods of research of body composition in athletes// *Электронный научный журнал «Биология и интегративная медицина» №4 – июль-август (44) 2020*– С.16-29
9. Rasulovna R. M. Sports Genetics is the Key to High Achievements of Athletes // *INTERNATIONAL JOURNAL OF HEALTH SYSTEMS AND MEDICAL SCIENCES*. – 2023. – Т. 2. – №. 1. – С. 23-30.
10. Rasulovna R. M. The Role of ADRB2, ADRB3 Genes Polymorphism in the Development of Age-Dependent Adaptability, Movement Speed, Speed-Strength Qualities in Junior and Cadet Athletes // *Scholastic: Journal of Natural and Medical Education*. – 2023. – Т. 2. – №. 1. – С. 147-152.
11. Рахматова М. Р., Жалолова В. З. Methods of research of body composition in athletes // *биология и интегративная медицина*. – 2020. – №. 4. – с. 16-28.
12. Гладков В. Н. Некоторые особенности заболеваний, травм, перенапряжений и их профилактика в спорте высших достижений / В. Н. Гладков. – М. : Сов. спорт, 2007. – 386 с.
13. Граевская Н. Д. Спортивная медицина: курс лекций и практические занятия / Н. Д. Граевская, Т. И. Довлатова. – М. : Сов. спорт, 2005. – 299 с.
14. Гурьянов М. С. Состояние здоровья и пути совершенствования медицинского обеспечения детско-юношеских спортивных школ : автореф. дис. канд. мед. наук / М. С. Гурьянов. – Казань, 2002. – 22 с.
15. Деревоедов В. В. Профессиональные заболевания в спорте высших достижений / В. В. Деревоедов. – М. : ЛФК и массаж, спортивная медицина. – 2008. – №8 (56). – С. 3–6.
16. Клейн К. В. Проблемы возрастных норм допуска к занятиям спортом детей и подростков / К. В. Клейн, И. В. Николаева, А. В. Люлюшин // *Материалы I Всероссийского конгресса «Медицина для спорта»*. – М., 2011. – С. 196–198.
17. Комолятова В. Н. Электрокардиографические особенности у юных элитных спортсменов / В. Н. Комо-лятова, Л. М. Макаров, В. О. Колосов, И. И. Киселева, Н. Н. Федина// *Педиатрия*. – 2013. – Т. 92, № 3. –С. 136–140.

18. Курникова М. В. Состояние морфофункционального статуса высококвалифицированных спортсменов подросткового возраста : автореф. дис. канд. мед. наук / М. В. Курникова. – М., 2009. – 22 с.
19. Мавлянов З.И., Жалолова В.З., Рахматова М.Р., Юлдашева Н.М. Характеристика компонентного состава гена FABP2 у юных спортсменов занимающихся различными видами спорта // Тиббиётда янги кун. – 2019. - № 4. – С. 35-42
20. Мавлянов З.И. Особенности соматотипа спортсмена и его взаимосвязь со спортивными генами. Дисс. Раб. на соиск. Учен. Степ. PhD. – 2018. – С. 18
21. Мавлянов З.И., Жалолова В.З., Рахматова М.Р., Анализ антропометрических показатели физического развития у юниоров и кадетов в спортивной медицине // Тиббиётда янги кун – 2020. - № 2(30/2). – С. 38-42
22. Мирошникова Ю. В. Медико-биологическое в обеспечение детско-юношеском спорте в Российской Федерации (концепция) / Ю. В. Мирошниченко, А. С. Самойлов, С. О. Ключникова, И. Т. Выходец // Педиатрия. – 2013. – Т. 92, № 1. – С. 143–149.
23. Михалюк Е. Л. Современные взгляды на диагностику метаболической кардиомиопатии вследствие хронического физического перенапряжения организма спортсменов / Е. Л. Михалюк, В. В. Сывовол // Спортивная медицина. – 2014. – № 1. – С. 3–12.
24. Ніколаєв С. Ю. Оздоровча спрямованість засобів атлетичної гімнастики для юнаків старшого шкільного віку / С. Ю. Ніколаєв // Молодіжний науковий вісник. – 2013. – № 9. – С. 85–88.
25. Платонов В. Н. Система подготовки спортсменов в олимпийском спорте. Общая теория и практические приложения// В. Н. Платонов. – Киев : Олимп. лит., 2004. – 808 с.
26. Расуловна, Р. М. . (2022) “Нейрофизиологический Статус Спортсменов Юниоров И Кадетов Занимающихся Легкой Атлетикой И Велоспортом”, *Miasto Przyszłości*, 25, pp. 217–220.
27. Рахматова М.Р., Жалолова В.З. Юниор ва кадет спортсменларда тананинг композицион таркибини ўрганиш.// Тиббиётда янги кун. - № 2 (30/2). - В. 67-
28. Рахматова М. Р., Собирова Г. Н. Спортчиларда Генлар Полиморфизмининг Ёшга Боғлиқ Мослашувчанлик, Ҳаракат Тезлиги, Тезлик-Куч Сифатлари Ривожланишидаги Аҳамияти //Miasto Przyszłości. – 2023. – Т. 36. – С. 266-271.