

## **Morphometric Parameters of Epididymis of Rats of Different Ages Receiving Radiation**

**Namozov Farrukh Jumaevich**

Bukhara State Medical Institute, Bukhara, Uzbekistan

**Abstract:** In this article, the morphometric parameters of the epididymis of rats of different ages that received radiation were examined. It has been proven in the experiment that the radiation, which has a negative effect on the morphological structures of the epididymis, has a significant effect on all the morphometric indicators of this organ. Chronic radiation sickness affects the vasculature of the epididymis and convoluted tubules, where the diameter of arterioles and venules, as well as the diameter of the convoluted tubules, is further reduced by radiation.

**Keywords:** epididymis, radiation, rat, morphology, morphometry.

**Relevance of the study.** Male infertility has many causes, ranging from unhealthy lifestyle choices to exposure to various chemical and physical factors. Among them, ionizing radiation has a special place, because it is classified as a group 1 carcinogen by the International Agency for Research on Cancer and is included in the list of occupational carcinogens by the International Labor Organization. Nevertheless, "the number of radiation facilities and the number of their employees are increasing by approximately 15 and 6% per year, respectively." This emphasizes once again that reproductive health and related aspects are not only a medical but also a social problem. All over the world, special attention is being paid to research aimed at improving the early detection, treatment and prevention of diseases of the male reproductive system, including the epididymis, caused by various physical factors. In this regard, determining the amount of ionizing radiation that causes pathological changes in the male body; development of a health monitoring system for workers working with radiation sources; determine the level of risk of infertility and the occurrence and development of tumors in these individuals; The study of the mechanisms of action of drugs that reduce radiation exposure remains a priority area of scientific research. Assessment of the morphofunctional state of the testicles after long-term exposure to ionizing radiation in chronic radiation sickness, comparison of the morphometric and morphological characteristics of these organs, and at the same time, development of an optimal algorithm for prevention, diagnosis and treatment are among the priority tasks for scientists. Targeted measures are being implemented in our country to fundamentally improve the healthcare system and improve the quality of medical services to the population. For this purpose, a number of tasks were defined, including "increasing the efficiency, quality and accessibility of medical care, implementing comprehensive measures aimed at supporting a healthy lifestyle and preventing diseases, including a healthy lifestyle the formation of a system of medical standardization, the introduction of high-tech diagnostic methods and treatment by creating effective models of patronage and clinical examination. Despite the positive results achieved in improving the quality of treatment of radiation-induced diseases of the reproductive system, in particular, oncological diseases, there is a need to develop new evidence-based treatment methods for specialists in this field. Solving the set tasks allows to reduce morbidity and mortality due to complications of oncological diseases that develop as a result of radiation,

raise the quality of modern medical services in the diagnosis and treatment of diseases to a new level, and improve the use of modern technologies in the provision of quality medical services.

**The purpose of the study** is to determine the morphometric parameters of the testicles of rats of different ages that received radiation.

**Material and methods.** An experimental study was conducted on the material obtained from the testicles of 100 purebred rats aged between 180 and 360 days, which were housed in a vivarium with a 12-hour light regime, access to standard food and water. At the beginning of the experiment, all sexually mature rats were quarantined for a week, and after the exclusion of somatic or infectious diseases, they were transferred to the usual vivarium regime. The animals were divided into 2 groups (n=100): Group I - rats that received radiation at a dose of 0.2 Gy (total dose 4.0 Gy) for 20 days from the age of 180 days (n=50); II - group - rats receiving radiation at a dose of 0.2 Gy (total dose 4.0 Gy) for 20 days from the age of 360 days (n=50).

The animals were slaughtered in the morning, on an empty stomach, under ether anesthesia by decapitation. After the abdominal cavity was opened, the extra testicles were removed and their mass, length, width, volume and tissue density were studied. The mass of each testicle was measured on an electric scale, and the length and width were measured with a millimeter tape. The obtained testicles were fixed in Bouin's solution. After passing through alcohol solutions of increasing concentration, they were embedded in hot paraffin, and then sections of testicular tissues with a standard thickness of 6-7  $\mu\text{m}$  were prepared, which were oriented sagittally or frontally. Sections were stained with hematoxylin and eosin according to Van Gieson. Prepared histological preparations were examined under a NLCD-307B binocular microscope (Novel, China).

**Research results.** In the 180-day-old male rats of the experimental group, the testicles were oval in shape and covered with a membrane whose thickness varied from 16.4 to 29.42  $\mu\text{m}$ , with an average of  $23.2 \pm 1.324 \mu\text{m}$ . Its wall is represented by bundles of collagen and elastic fibers. Bundles of elastic fibers are thin and mainly oriented longitudinally. In places, thin reticular fibers form a fine ring network. The outer surface of the shell is covered with 2-3 layers of longitudinally oriented flat mesothelial cells. In the 180-day-old male rats of the experimental group, the testicles were oval in shape and covered with a membrane whose thickness varied from 16.4 to 29.42  $\mu\text{m}$ , with an average of  $23.2 \pm 1.324 \mu\text{m}$ . Its wall is represented by bundles of collagen and elastic fibers. Bundles of elastic fibers are thin and mainly oriented longitudinally. In places, thin reticular fibers form a fine ring network. The outer surface of the shell is covered with 2-3 layers of longitudinally oriented flat mesothelial cells. Under the testicular membrane there are intrathecal vessels: arterioles with sizes from 127.94x127.94 to 132.43x132.43  $\mu\text{m}$ , the average is  $130.6 \pm 0.433 \mu\text{m}$ . Their wall thickness varies from 18.03 to 32.04  $\mu\text{m}$ , with an average of  $25.1 \pm 1.459 \mu\text{m}$ . These arterioles pass in a transverse direction from the front, free edge of the testis to the surface where the epididymis is located. The distance between arterioles is from 360 to 775  $\mu\text{m}$ , and their number is 9-12. The diameter of the venules is from 23.64x33.85 to 36.41x53.71  $\mu\text{m}$ , the average is  $50.3 \pm 0.865 \mu\text{m}$ , the wall thickness is  $5.4 \pm 0.259 \mu\text{m}$ .

Convolutated seminiferous tubules form the base of the testis. Diameter - 144.52 - 192.22  $\mu\text{m}$  (average -  $172.5 \pm 4.014 \mu\text{m}$ ). The cross-sectional area of the tubes is 20198.4 - 27443.6  $\mu\text{m}^2$  (average -  $23358.6 \pm 782.48 \mu\text{m}^2$ ). (see Fig. 1).

The basis of the tubular wall is the fibrous structure of connective tissue, which consists of reticular fibers and bundles of collagen and elastic fibers. Reticular fibers around the seminiferous tubules form a "honeycomb"-like network in transverse histological sections. The thickness of the spermatogenic epithelium ranges from 43.45 to 68.99  $\mu\text{m}$ , the average is  $53.1 \pm 2.283 \mu\text{m}$ .

The first, more strongly stained layer of cells is the Spermatagonium. Their average size is  $8.5 \pm 0.327 \mu\text{m}$ . The second layer consists of larger cells - spermatocytes (size  $12.5 \pm 0.148 \mu\text{m}$ ).



**Fig. 1. Testicles of a 180-day-old rat suffering from chronic radiation sickness. 1- convoluted seed tubules; 2-channel space. Stained with hematoxylin-eosin. Ob. 10 x Ok. 10**

The third layer of spermatids, their average size is  $8.3 \pm 0.120 \mu\text{m}$ . The diameter of the free space (light zone) of the seminiferous tubule is  $42.6\text{--}61.5 \mu\text{m}$ , on average -  $55.6 \pm 2.13 \mu\text{m}$ . Accompanying arterioles, venules, and capillaries are located in the intercanal spaces. The dimensions of intertubular arterioles range from  $29.71 \times 29.71$  to  $33.47 \times 33.47 \mu\text{m}$ , the average is  $31.5 \pm 0.322 \mu\text{m}$ , the average wall thickness is  $9.9 \pm 0.193 \mu\text{m}$ . The inner membrane of arterioles is formed from endothelial cells with elongated nuclei located in the basement membrane. Outside the inner shell, there is an internal elastic membrane, which has a well-defined fold. The middle shell consists of a series of spirally arranged smooth muscle cells, the nuclei of which are spindle-shaped. The outer shell is made of loose connective tissue. The wall of intertubular venules is formed from one layer of endothelocytes. The average wall thickness of venules is  $5.3 \pm 0.220 \mu\text{m}$ , and the cavity diameter is  $27.3 \pm 0.721 \mu\text{m}$ . The capillary cavity diameter is on average  $5.8 \pm 0.378 \mu\text{m}$ . Their wall thickness is  $3.9 \pm 0.225 \mu\text{m}$ . Leydig cells - testicular endocrinocytes - are located in the interductal spaces in a triangular shape. The size of Leydig cells is from  $7.43$  to  $10.61 \mu\text{m}$ , the average is  $9.2 \pm 0.290 \mu\text{m}$ , and their number in the field of vision is from  $25$  to  $39$ , the average is  $31.8 \pm 1.269$ . Reticular and collagen fibers in the intertubular spaces form a network, in the rings of which Leydig cells are located.

In the 360-day-old male rats in the experimental group, the testicles have an oval shape, are covered with a protein film on the outside, and its frame is made of bundles of reticular fibers and collagen fibers. In some places bundles of elastic fibers are identified. Outside, it is covered with 2-4 rows of flat mesothelial cells. The shell wall thickness ranges from  $20.2$  to  $38.05 \mu\text{m}$ , with an average of  $29.4 \pm 1.301 \mu\text{m}$ . Under the testicular capsule there are subcapsular arterioles with diameters ranging from  $148.66 \times 148.66$  to  $155.54 \times 155.54 \mu\text{m}$ , with an average of  $152.3 \pm 0.577 \mu\text{m}$ . The average wall thickness of arterioles is  $33.7 \pm 0.822 \mu\text{m}$ . Arterioles are located at a distance of  $374$  to  $796 \mu\text{m}$ . In addition to them, venules are identified under the capsule, their sizes range from  $29.85 \times 42.61$  to  $55.14 \times 72.40 \mu\text{m}$ , the average is  $69.1 \pm 0.525 \mu\text{m}$ . Their wall thickness is  $6.0 \pm 0.351 \mu\text{m}$ . The diameter of the testis is from  $205.96$  to  $227.57 \mu\text{m}$ , on average -  $216.2 \pm 1.976 \mu\text{m}$ , based on curved seminiferous tubules. The field of vision of seminiferous tubules ranges from  $34321.7$  to  $40054.6 \mu\text{m}^2$ , the average is  $36692.8 \pm 573.29 \mu\text{m}^2$ . Spermatogenic epithelium is located inside the wall of seminiferous tubules. Its height ranges from  $57.51$  to  $73.46 \mu\text{m}$ , with an average of  $67.4 \pm 1.225 \mu\text{m}$ . The layered arrangement of the spermatogenic epithelium does not differ much from previous ages. In the cavity of the seminiferous tubules, there is an intercellular trophic substance that forms the light zone. The diameter of the free space is from  $60.2$  to  $78.7 \mu\text{m}$ , the average is  $69.8 \pm 1.85 \mu\text{m}$ .



**Fig. 2.** Testicles of a 360-day-old rat suffering from chronic radiation sickness. 1- convoluted seed tubules; 2-channel space. Stained with hematoxylin-eosin. Ob. 10 x Ok. 20

In the intertubular spaces there are Leydig cells, the size of which is 8.54-12.17  $\mu\text{m}$ . Average -  $11.0 \pm 0.310 \mu\text{m}$ . The number of Leydig cells ranges from 30 to 47, on average -  $41.3 \pm 1,230 \mu\text{m}$ . In addition, accompanying arterioles and venules are located in the intertubular spaces. The diameter of arterioles ranges from  $34.19 \times 34.19$  to  $37.24 \times 37.24 \mu\text{m}$ , with an average of  $35.8 \pm 0.265 \mu\text{m}$ . The average wall thickness of intertubular arterioles is  $12.2 \pm 0.302 \mu\text{m}$ . It consists of 3 shells. Internal - forms endothelial cells with elongated nuclei located in the basement membrane. Outside of it there is an internal elastic membrane, which has a well-defined tooth. The middle shell is represented by a single layer of spirally arranged smooth muscle cells with elongated nuclei. The outer shell is made of loose connective tissue. In the wall of intertubular venules there are single-layered endothelial cells with large nuclei. The average wall thickness of venules is  $6.5 \pm 0.382 \mu\text{m}$ . The average diameter of the capillary cavity is  $6.9 \pm 0.410 \mu\text{m}$ . The capillary wall is formed by a single layer of endothelocytes with an elongated nucleus located at a close distance from each other. The wall thickness of the capillaries is  $4.7 \pm 0.094 \mu\text{m}$ . Reticular fibers in the intertubular spaces form a fine ring network, surrounding the intertubular vessels and Leydig cells. Bundles of collagen fibers are multidirectional.

**Summary.** Irradiation, which has a negative effect on the morphological structures of the epididymis, leads to a delay in the development of spermatogenic epithelium in all morphometric parameters, including the thickness of the albuginea. Latency is more pronounced in purebred rats at 180 days and less at 360 days. This is related to the activation of the body's protective and compensatory mechanisms. Chronic radiation sickness affects the vasculature of the testes and convoluted tubules, where the diameter of arterioles and venules, as well as the diameter of the convoluted tubules, is further reduced by radiation.

#### REFERENCES:

1. Djumayevich N. F. Macro-Microscopic Anatomy and Reactive Changes in the Testimonial Appendices under Exposure to Magnesium Chlorate in Postnatal Ontogenesis //International Journal of Innovative Analyses and Emerging Technology. – 2021. – T. 1. – №. 6. – C. 152-155.
2. Jumaevich N. F. Comparative characteristics of testicular appendages in normal conditions and when exposed to a biostimulator against the background of radiation sickness //International journal for innovative engineering and management research. – 2021. – C. 28-30.

3. Samiyevich S. E., INTERVENTIONS N. F. J. E. OZONE THERAPY IN THE COMPLEX TREATMENT OF PATIENTS WITH MECHANICAL JAUNDICE AND CHOLANGITIS WITH CHOLEDOCHOLITHIASIS //ResearchJet Journal of Analysis and Inventions. – 2021. – Т. 9. – №. 2. – С. 22-27.
4. Турдиев М. Р., Махмудова Г. Ф. Морфофункциональные изменения, происходящие в селезенке в результате действия внешних и внутренних факторов //Тиббиётда янги кун. – 2022. – №. 11. – С. 49.
5. Turdiyev M. R., Sokhibova Z. R. Morphometric characteristics of the Spleen of white rats in normal and in chronic Radiation Disease //The american journal of medical sciences and pharmaceutical research. – 2021. – Т. 3. – №. 02. – С. 146-154.
6. Turdiev M. R., Teshayev S. J. Comparative characteristics of the spleen of white rats in normal and chronic radiation sickness //Chief Editor. – Т. 7. – №. 11.
7. Turdiyev M. R. Teshayev Sh //J. Morphometric Assessment of Functional Immunomorphology of White Rat Spleen in the Age Aspect American Journal of Medicine and Medical Sciences. – 2019. – Т. 9. – №. 12. – С. 523-526.
8. Турдиев М. Р. и др. ЧАСТОТА РАСПРОСТРАНЕНИЯ РАКА МОЛОЧНОЙ ЖЕЛЕЗЫ В БУХАРСКОЙ ОБЛАСТИ //Молодежный инновационный вестник. – 2015. – Т. 4. – №. 1. – С. 267-268.
9. Turdiev M. R. Teshayev Sh. J. Comparative characteristics of the morphological and morphometric parameters of the spleen of white rats in normal conditions, chronic radiation sickness and correction with a biostimulant //Problems of biology and Medicine. – 2020. – №. 4. – С. 120.
10. Турдиев М. Р., Сохибова З. Р. Этиологические факторы острых аллергических состояний у детей, проживающих в условиях города Бухары //Новый день в медицины. – 2018. – №. 3. – С. 23.
11. Турдиев М. Р. Морфофункциональные особенности селезенки белых крыс в норме и при хронической лучевой болезни //Новый день в медицине.–2020.–3 (31)–С. – С. 734-737.
12. Turdiyev M. R., Sanoyev B. A. Pathologi of the afterbirth during 2020 in the Bukhara regional perinatal center //Eurasian Journal of Medical and Natural sciences. Volume1. – 2021. – №. 2.
13. Turdiev M. R. et al. ChASTOTA RASPROSTRANENIYa RAKA MOLOChNOY ZhELEZY V BUKhARSKOY OBLASTI //Молодежный инновационный вестник. – 2015. – Т. 4. – №. 1. – С. 267-268.
14. Turdiev M. R. Morphological and morphometric parameters of lymphoid Structures of the Spleen of white rats in Postnatal ontogenesis in Dynamics of Age. European multidisciplinary journal of modern science. Volume 4, 2022. – P-319-326.
15. Turdiyev M. R. Morphological and Orthometric Parameters of lymphoid Structures of the Spleen of white rats //Central Asian Journal of Medical and Natural Sciences. Volume. – Т. 2.
16. Turdiyev M. R. Morphometric Indicators of Morphological Structures of the White Rats Spleen in Postnatal Ontogenesis //Web of Synergy: International Interdisciplinary Research Journal. – 2023. – Т. 2. – №. 4. – С. 576-580.
17. Turdiyev M. R., Boboeva R. R. CHOLERETIC ACTIVITY OF RUTANA AT THERAPEUTIC APPLICATION IN RATS WITH HELIOTRIN HEPATITIS //Oriental renaissance: Innovative, educational, natural and social sciences. – 2021. – Т. 1. – №. 8. – С. 644-653.

18. Турдиев М. Р. Морфофункциональные Изменения Лимфоидных Структур Селезенки Белых Крыс В Постнатальном Онтогенезе В Динамике Возраста //AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI. – 2023. – Т. 2. – №. 5. – С. 188-192.
19. Turdiev M. R. Morphofunctional Changes in Lymphoid Structures of the Spleen of White Rats in Postnatal Ontogenesis in the Dynamics of Age //Web of Synergy: International Interdisciplinary Research Journal. – 2023. – Т. 2. – №. 5. – С. 144-148.
20. Rustamovich T. M. et al. Edematous Breast Cancer Problems of Diagnosis and Treatment //Research Journal of Trauma and Disability Studies. – 2022. – Т. 1. – №. 10. – С. 93-100.
21. Turdiev M. R. Histological Analysis of the Spleen of White Rats in Postnatal Ontogenesis //Research Journal of Trauma and Disability Studies. – 2022. – Т. 1. – №. 10. – С. 135-141.
22. Rustamovich T. M. Morphological and Orthometric Parameters of Lymphoid Structures of the Spleen of White Rats //Central Asian Journal of Medical and Natural Science. – 2021. – Т. 2. – №. 5. – С. 122-128.
23. Олимова А. З., Шодиев У. М. Репродуктив Ёшдаги эркакларда бепуштлик сабаблари: Бухоро тумани эпидемиологияси //Scientific progress. – 2021. – Т. 2. – №. 7. – С. 499-502.
24. Zokirovna O. A., Abdurasulovich S. B. Ovarian Diseases in Age of Reproductive Women: Dermoid Cyst //IJTIMOIY FANLARDA INNOVASIYA ONLAYN ILMIY JURNALI. – 2021. – Т. 1. – №. 6. – С. 154-161.
25. Olimova A. Z. ECHINOCOCCOSIS OF LIVER OF THREE MONTHLY WHITE RAT //Scientific progress. – 2022. – Т. 3. – №. 3. – С. 462-466.
26. Олимова А. З. Морфологические и морфометрические особенности печени белых беспородных трех месячных крыс после тяжелой черепно-мозговой травмы вызванной экспериментальным путём //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2021. – Т. 1. – №. 6. – С. 557-563.
27. Oglu M. Z. M., Zokirovna O. A. МОРФОЛОГИЧЕСКИЕ И МОРФОМЕТРИЧЕСКИЕ ПАРАМЕТРЫ ПЕЧЕНИ БЕЛЫХ БЕСПОРОДНЫХ КРЫС, ПЕРЕНЕСШИХ ЭКСПЕРИМЕНТАЛЬНУЮ ЧЕРЕПНО-МОЗГОВУЮ ТРАВМУ ПОСЛЕ МЕДИКАМЕНТОЗНОЙ КОРРЕКЦИИ //JOURNAL OF BIOMEDICINE AND PRACTICE. – 2023. – Т. 8. – №. 1.
28. Олимова А. З., Турдиев М. Р. БУХОРО ШАХРИДА МЕЪДА ВА ЁН ИККИ БАРМОҚЛИ ИЧАК ЯРАСИ УЧРАШ ЭПИДЕМИОЛОГИЯСИ //Oriental renaissance: Innovative, educational, natural and social sciences. – 2022. – Т. 2. – №. 4. – С. 642-647.
29. Zokirovna O. A. Modern Concepts of Idiopathic Pulmonary Fibrosis //American Journal of Pediatric Medicine and Health Sciences. – 2023. – Т. 1. – №. 3. – С. 97-101.
30. Zokirovna O. A. Pathology of Precancerous Conditions of the Ovaries //American Journal of Pediatric Medicine and Health Sciences. – 2023. – Т. 1. – №. 3. – С. 93-96.
31. Зокировна, Олимова Азиза и Тешаев Шухрат Джумаевич. «Морфологические аспекты печени белых беспородных крыс после тяжелой черепно-мозговой травмы, вызванной экспериментально в виде дорожно-транспортного происшествия». *Scholastic: Journal of Natural and Medical Education* 2.2 (2023): 59-62.
32. Zokirovna O. A. Comparative characteristics of the morphological parameters of the liver at different periods of traumatic brain injury //Euro-Asia Conferences. – 2021. – С. 139-142.
33. Zokirovna O. A. Macroand microscopic structure of the liver of threemonthly white rats //Academic research in educational sciences. – 2021. – Т. 2. – №. 9. – С. 309-312.
34. Олимова А. З. Частота Встречаемости Миомы Матки У Женщин В Репродуктивном Возрасте //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2021. – Т. 1. – №. 6. – С. 551-556.

35. Zokirovna O. A., Abdurasulovich S. B. Ovarian Diseases in Age of Reproductive Women: Dermoid Cyst //IJTIMOIY FANLARDA INNOVASIYA ONLAYN ILMIY JURNALI. – 2021. – T. 1. – №. 6. – C. 154-161.
36. Zokirovna O. A. Cytological screening of cervical diseases: pap test research in the bukhara regional diagnostic center for the period 2015-2019. – 2022.
37. Zokirovna O. A., PREVALENCE R. M. M. EPIDEMIOLOGY OF CANCER OF THE ORAL CAVITY AND THROAT IN THE BUKHARA REGION //Web of Scientist: International Scientific Research Journal. – 2022. – T. 3. – №. 11. – C. 545-550.
38. Olimova A. Z. The frequency of occurrence of my uterus In women of reproductive age //JOURNAL OF ADVANCED RESEARCH AND STABILITY (JARS). – 2021. – T. 1. – №. 06. – C. 551-556.
39. Olimova Aziza Zokirovna. (2023). MODERN PRINCIPLES OF THE EFFECT OF HEMODIALYSIS THERAPY ON HEART RATE. *International Journal of Integrative and Modern Medicine*, 1(1), 80–85. Retrieved from <http://medicaljournals.eu/index.php/IJIMM/article/view/28>
40. Olimova Aziza Zokirovna. (2023). PATHOMORPHOLOGICAL CHARACTERISTICS OF THE EPIDIDYMISS UNDER IRRADIATION. *International Journal of Integrative and Modern Medicine*, 1(1), 96–100. Retrieved from <http://medicaljournals.eu/index.php/IJIMM/article/view/31>
41. Olimova Aziza Zokirovna. (2023). THE INCIDENCE OF CANCER OF THE ORAL CAVITY AND PHARYNX IN THE BUKHARA REGION. *International Journal of Integrative and Modern Medicine*, 1(1), 86–89. Retrieved from <http://medicaljournals.eu/index.php/IJIMM/article/view/29>
42. Olimova Aziza Zokirovna. (2023). INFLUENCE OF ALCOHOL INTOXICATION ON THE HEART TISSUE OF RATS IN THE EXPERIMENT. *International Journal of Integrative and Modern Medicine*, 1(1), 90–95. Retrieved from <http://medicaljournals.eu/index.php/IJIMM/article/view/30>
43. Olimova Aziza Zokirovna. (2023). Modern Aspects of the Etiology of Gastric Ulcer and Its Complications. *American Journal of Pediatric Medicine and Health Sciences (2993-2149)*, 1(3), 163–166. Retrieved from <http://grnjournal.us/index.php/AJPMHS/article/view/208>
44. Zokirovna O. A., Jumaevich T. S. Morphological Aspects of the Liver of White Outbred Rats After Severe Traumatic Brain Injury Caused Experimentally in the Form of a Road Accident //Scholastic: Journal of Natural and Medical Education. – 2023. – T. 2. – №. 2. – C. 59-62.
45. Aziza Zokirovna Olimova GASTRIC ULCER AND ITS COMPLICATIONS // Scientific progress. 2022. №3. URL: <https://cyberleninka.ru/article/n/gastric-ulcer-and-its-complications> (дата обращения: 28.09.2023).
46. Olimova Aziza Zokirovna. (2022). TECHNIQUE FOR CUTTING BIOPSY AND SURGICAL MATERIAL IN THE PRACTICE OF PATHOLOGICAL ANATOMY AND FORENSIC MEDICINE. *Web of Scientist: International Scientific Research Journal*, 3(7), 116–120. <https://doi.org/10.17605/OSF.IO/PSQ59>
47. Zhumayevich N. F., Zokirovna O. A. PATHOMORPHOLOGY OF GASTRIC CANCER //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2022. – C. 330-333.
48. Zokirovna O. A. Epidemiological and Etiological Data of Morphogenesis and Pathomorphology of Congenital Heart Diseases in Children //American Journal of Pediatric Medicine and Health Sciences. – 2023. – T. 1. – №. 4. – C. 88-91.