

Estrogen (Er- A) And Progesterone (Pr- A) Receptors Expression In Egg Endometriosis

Juraeva Gulbakhor Baxshillaevna

Associate Professor, Candidate of Medical Sciences, Head of the Department of Pathological Anatomy, Bukhara, Uzbekistan. E-mail: gjurayeva20@gmail.com

Annotation: This paper examines the degree of expression of estrogen (ER- α) and progesterone (PR- α) receptors in both glandular epithelium and stromal structures in ovarian endometriosis. As a material, ovarian tissue from 28 patients was immunohistochemically examined. Immunohistochemical examination of ovarian endometriosis revealed differences in the expression of estrogen (ER- α) and progesterone (PR- α) receptors in both epithelial cells and stromal structures. The sensitivity of these tissue structures to hormones during endometriosis, and the formation of specific protein receptors indicates the high rate of expression of the ER- α receptor relative to the PR- α receptor in epithelial cells that have become endometriotic glands relative to stroma structures indicates that the role of the hormone estrogen in the mechanism of endometriosis development is highly important.

Keywords: ovary, endometriosis, immunohistochemistry, estrogen, progesterone, ER-a, PR-a receptors.

INTRODUCTION

In the study of endometriosis by morphological, histochemical, and immunohistochemical methods, it is important to know the histogenesis of the origin of the endometrial glands and epithelial glands of organs and tissues where endometriosis develops. After all, the metaplastic theory of the development of endometriosis plays a key role. The ovarian and Mueller tubes of the ovary arise from the selemic mesothelium, so the germinative epithelium of the ovary becomes the endometrial glands. However, because the mesothelium of the peritoneal serous membrane is a multipotent tissue, abdominal endometriosis develops from in-situ metaplasia of the mesothelium (1, 2).

In the study of the pathogenesis of endometriosis, Sampson advanced the theory of implantation. Because menstrual blood contains endometrial cells, they are implanted in the abdomen and other organs, proliferate, and lead to endometriosis. Hence, retrograde shortening of the fallopian tube causes retrograde menstruation and is a confirmation of the development of endometriosis in the fallopian tube and abdominal cavity (3, 4, 5). It is estimated that the dissemination of endometrial cells through lymph and blood vessels causes extragenital forms of endometriosis.

Sex hormones play a key role in the pathogenesis of endometriosis. Several clinical and experimental studies have confirmed that endometriosis is an estrogen-dependent disease. As evidence of this, endometriosis does not develop in girls until menopause begins. Endometriosis is often confirmed to develop due to high levels of estrogen, estrogenic obesity, and the artificial introduction of estrogen medication.

Immunohistochemical examination detects protein-type receptors located on the surface of estrogen and progesterone-sensitive cells, i.e., antigen-specific antibodies. Because the epithelium of other organs and tissues outside the uterus originates from the selemic epithelium, receptors sensitive to specific estrogen and progesterone hormones appear, affecting them as the number of sex hormones in the body increases (3,4). Detection of these receptors by antigen-antibody reaction in the immunohistochemical examination, their positive staining with a specially marked antibody confirms the presence of these receptors, the development of endometriosis. The results of this immunohistochemical method are evaluated by the following accepted terms: "strong positive reaction", "false positive reaction", "negative reaction" and "false negative reaction".

MATERIALS AND METHODS

As a material, a total of 28 patients with an average age of 28.5 ± 4.2 years who underwent surgery in the RPAC of the Republic of Uzbekistan biopsy diagnostics department in 2016-2021 underwent macroscopic examination of ovaries surgically removed with a diagnosis of endometriosis and incisions were made from areas where endometriosis-specific glands and cysts had grown. Biopsy sections were incubated for 48h in 10% neutralized formalin. Dehydration was carried out at increasing concentrations of alcohols and chloroform. Histological incisions were initially stained with hematoxylin and eosin to determine topography. Then a series of cuts from paraffin bricks were carried out in a specially automated system Ventana Benchmark XT, Roche, Switzerland, deparaffinization, dehydration, demasking, and staining in antigens. Receptors sensitive to estrogen (ER- α) and progesterone (PR- α) were identified using specially labeled antibodies. These receptors for weak, moderate, and strongly positively stained glandular epithelium and stromal cells were calculated by the following formula: $HS = 1a + 2b + 3c$, where a is the percentage of weakly stained cells, b is the percentage of moderately stained cells, and s is the percentage of strongly stained cells. The results were calculated on the following points: 0-10 points - no expression, 11 - 80 points - slow expression, 81 - 140 points - average expression, 141 - 300 points - strong expression. Statistical analysis of the obtained quantitative indicators is a practical statistical analysis program (Statistica Windows 10.0, SPSS Statistics 22, Microsoft Excel). In the groups, the performance of the characters was assessed using the Manna-Whitney test, and the statistical evaluation of the characters was performed at $p \leq 0.05$.

RESULTS AND DISCUSSION

The results of the study were determined as a percentage of slow, moderate, and high levels of estrogen (ER- α) and progesterone (PR- α) receptor expression. Estrogen (ER- α) receptor expression in the ovarian glandular epithelium was strong in 72.6% of 28 patients - an average of 286.4 points, 19.2% - 138.6 points, and 8.2% - 56.8 points. expressed. In cases of strong expression of the estrogen (ER-a) receptor, the epithelium of endometriotic gland structures growing under the outer mesothelium of the ovary is located in 2-3 rows, and both the cytoplasm and the nucleus of the ER-a receptor are strongly brown (Fig. 1). In the single-layered prismatic epithelium covering the outer surface of the ovary, it is observed that this receptor is almost not expressed. ER- α receptors are also relatively well expressed in the cells of the stroma-vascular structures around the gland.

In 8.2% of cases of ovarian endometriosis, the epithelium of endometriotic glandular structures consists mainly of single-layered prismatic and cylindrical epithelium, and in their cytoplasm, the ER-a receptor is expressed in a slightly light brown form (Fig. 2). In the epithelial nuclei of some of these glands, this receptor is found to be relatively well expressed. It is also found that this receptor is less expressed in the stroma cells around the glands than in the previous group of patients.

In cases of ovarian endometriosis, the level of estrogen (ER- α) receptor expression in the cells of stroma-vascular structures was also different, i.e. strong expression in 56.8% (264.8 points), moderate expression in 28.7% (134.6 points) and slow expression in 14.5% (48.6 points). Studies of ovarian stroma structures have shown that the ER- α receptor is well expressed in the cytoplasm of most histiocytic cells, as well as in the nuclei of some. Most of the histiocytic cells are large, and in their cytoplasm, it is found that this receptor is expressed in the form of granules of various sizes from dark brown to light golden (Fig. 3). Depending on the degree of maturation of the cells, i.e. in less differentiated histiocytic cells, it is observed that this receptor is less expressed. In some very small and lymphoid-like structures, it is found to be in the form of light golden granules resembling tiny bubbles.




Figure 1. Strong expression of the ER- α receptor in ovarian endometriotic glands. Stain: immunohistochemical method. X400.




Figure 2. Slow expression of the ER- α receptor in ovarian endometriosis glands. Stain: immunohistochemical method. X400.




Figure 3. Ovarian endometriosis, good expression of ER- α receptor in stroma cells. Stain: immunohistochemistry. X450.




Figure 4. Ovarian endometriosis, slow expression of the ER- α receptor in stroma cells. Stain: immunohistochemistry. X650.

Determination of the degree of expression of the progesterone (PR- α) receptor in ovarian endometriosis showed that this receptor was also expressed at different levels in the glandular cells and stroma structures. Progesterone (PR- α) receptor expression in the ovarian glandular epithelium was strong in 62.6% of 28 patients - an average of 262.4 points, 25.3% - 128.2 points and 12.1% - 36.8 points expressed. Morphologically, the expression of the progesterone (PR- α) receptor has specific characteristics. When this receptor was strongly expressed in the glandular epithelium, it was observed that the nuclei of the cell were mostly stained dark brown (Fig. 5). Another peculiarity was that this receptor was found to be expressed only in stromal cells around the glands.

Low expression of this receptor was observed at some relatively strong levels of glandular epithelium and very low levels at others (Fig. 6). In the stromal structures surrounding such glands, however, expression was not observed only in individual cells and in most histiocytic cells.




Figure 5. Ovarian endometriosis, expression of progesterone (PR-α) receptors in both epithelial and stromal cells. Stain: immunohistochemistry. X400




Figure 6. Ovarian endometriosis, expression of progesterone (PR-α) receptors in both epithelial and stromal cells. Stain: immunohistochemistry. X400

CONCLUSION

Immunohistochemically examination of ovarian endometriosis revealed differences in the expression of estrogen (ER-α) and progesterone (PR-α) receptors in both epithelial cells and stromal structures. The sensitivity of these tissue structures to hormones during endometriosis, and the formation of specific protein receptors indicates the high rate of expression of the ER-α receptor relative to the PR-α receptor in epithelial cells that have become endometriosis glands relative to the stroma structures indicates that the role of the hormone estrogen in the mechanism of endometriosis development is highly important. A deeper study of the composition of these receptors at the level of molecular biology will help to determine what macromolecular reactions underlie the pathogenesis of endometriosis.

LIST OF REFERENCES

1. BA S., Israilov R. I., Djuraeva G. B. Quantitative indicators and methods for modeling structural units in placental insufficiency //World Journal of Pharmaceutical Research. – 2020. – Т. 9. – №. 12. – С. 37-47.
2. Samievich S. E. Jurayeva Gulbaxor Bakhshilloeyvna Bronchopulmonary Complications After Heart Surgery With Congenital Defects //International Journal For Innovative Enjineering And Management Rewsearch. – Т. 10. – С. 320-323.
3. Жураева Г. Б., Саидов А. А., Исраилов Р. И. Зависимость морфологических и морфометрических изменений слизистой оболочки при острой кишечной инфекции от вида возбудителя //Молодежный инновационный вестник. – 2015. – Т. 4. – №. 1. – С. 193-195.
4. Bakhshillaevna Z. G. Pathomorphological Characteristics Of Glandular Hyperplasia Endometry In Women According To The Data Of The Bukhara Regional Pathological Bureau //The American Journal of Interdisciplinary Innovations and Research. – 2020. – Т. 2. – №. 11. – С. 142-149.
5. Жураева , Г. (2021). ИЗУЧЕНИЕ КЛИНИКО-МОРФОЛОГИЧЕСКИХ ОСОБЕННОСТЕЙ РАЗНЫХ ФОРМ ЭНДОМЕТРИОЗА. *Збірник наукових праць SCIENTIA*. вилучено із <https://ojs.ukrlogos.in.ua/index.php/scientia/article/view/9992>

6. Bakhshillaevna J. G. CHARACTERISTICS AND ROLE OF ESTROGEN (ER-A) AND PROGESTERONE (PR-A) IN OVARIAN ENDOMETRIOSIS //International Conference on Research Identity, Value and Ethics. – 2022. – С. 194-198.
7. BA S., Israilov R. I., Djuraeva G. B. Quantitative indicators and methods for modeling structural units in placental insufficiency //World Journal of Pharmaceutical Research. – 2020. – Т. 9. – №. 12. – С. 37-47.
8. Bakhshillaevna Z. G. Pathomorphological Characteristics Of Glandular Hyperplasia Endometry In Women According To The Data Of The Bukhara Regional Pathological Bureau //The American Journal of Interdisciplinary Innovations and Research. – 2020. – Т. 2. – №. 11. – С. 142-149.
9. Gulbakhor J. Study of clinical and morphological features of different form of endometrioid disease //Жураева Г.Б.//Общество и инновации: М. – 2021. – Т. 5. – С. 265-270.
10. Жураева Г. Б. и др. Корреляционная взаимосвязь между клинико-лабораторными показателями и структурными изменениями в почках при каждой морфологической форме хронического гломерулонефрита, сопровождающейся нефротическим синдромом //Вестник Совета молодых учёных и специалистов Челябинской области. – 2017. – Т. 3. – №. 2 (17). – С. 33-35.
11. Жураева Г.Б., Саидов А.А., Исраилов Р.И. ЗАВИСИМОСТЬ МОРФОЛОГИЧЕСКИХ И МОРФОМЕТРИЧЕСКИХ ИЗМЕНЕНИЙ СЛИЗИСТОЙ ОБОЛОЧКИ ПРИ ОСТРОЙ КИШЕЧНОЙ ИНФЕКЦИИ ОТ ВИДА ВОЗБУДИТЕЛЯ // Молодежный инновационный вестник. - 2015. - Т. 4. - №1. - С. 193-195.
12. Жураева Г. Б., Зикирова А. И. БАЧАДОНГА ҚЎШНИ АЪЗОЛАР ЭНДОМЕТРИОЗИДА ИММУНОГИСТОКИМЁВИЙ МАРКЕР КИ-67 ЭКСПРЕССИЯЛАНИШ ДАРАЖАСИ ВА ПРОЛИФЕРАТИВ ИНДЕКСИ //Oriental renaissance: Innovative, educational, natural and social sciences. – 2022. – Т. 2. – №. 10. – С. 156-164.
13. Жураева Г. Б. Гистопатология миометрия при внутреннем эндометриозе //Scientific progress. – 2022. – Т. 3. – №. 2. – С. 169-174.
14. Juraeva G. B., Zikirova A. I. Pathomorphological Changes Of Endometriosis In Women Of Bukhara Region //International Conference on Scientific Research and Advancements in Sciences. – 2021. – С. 134-136.
15. Bakhillaevna J. G., Ibodillaevna Z. A., Farkhodovna K. T. Morphological ogikal and cayses of dysfunctional uretine bleding //World bulletin of public healthn. – 2021. – Т. 5. – №. 12. – С. 38-41.
16. Жураева Г. Б. Иммуногистохимическое Исследования Эндометриоза Различных Органов С Помощью Маркера Ки-67 //AMALIY VA TIBBIYOT FANLARI ILMIY JURNALI. – 2022. – Т. 1. – №. 6. – С. 27-33.
17. Juraeva G. Competent factors for developing students' creative abilities in higher educational institutions //AIP Conference Proceedings. – AIP Publishing LLC, 2022. – Т. 2432. – №. 1. – С. 060028.
18. Bakhshillaevna Z. G. The Course Of Pathological Anatomy In A Modular System, As A New Teaching Technology In A Medical University //The American Journal of Applied sciences. – 2020. – Т. 2. – №. 11. – С. 122-126.

19. Олимова А. З., Шодиев У. М. Репродуктив Ёшдаги эркакларда бепуштлик сабаблари: Бухоро тумани эпидемиологияси //Scientific progress. – 2021. – Т. 2. – №. 7. – С. 499-502.
20. Zokirovna O. A., Abdurasulovich S. B. Ovarian Diseases in Age of Reproductive Women: Dermoid Cyst //IJTIMOIY FANLARDA INNOVASIYA ONLAYN ILMIY JURNALI. – 2021. – Т. 1. – №. 6. – С. 154-161.
21. Olimova A. Z. ECHINOCOCCOSIS OF LIVER OF THREE MONTHLY WHITE RAT //Scientific progress. – 2022. – Т. 3. – №. 3. – С. 462-466.
22. Олимова А. З. Морфологические и морфометрические особенности печени белых беспородных трех месячных крыс после тяжелой черепно-мозговой травмы вызванной экспериментальным путём //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2021. – Т. 1. – №. 6. – С. 557-563.
23. Oglu M. Z. M., Zokirovna O. A. МОРФОЛОГИЧЕСКИЕ И МОРФОМЕТРИЧЕСКИЕ ПАРАМЕТРЫ ПЕЧЕНИ БЕЛЫХ БЕСПОРОДНЫХ КРЫС, ПЕРЕНЕСШИХ ЭКСПЕРИМЕНТАЛЬНУЮ ЧЕРЕПНО-МОЗГОВУЮ ТРАВМУ ПОСЛЕ МЕДИКАМЕНТОЗНОЙ КОРРЕКЦИИ //JOURNAL OF BIOMEDICINE AND PRACTICE. – 2023. – Т. 8. – №. 1.
24. Олимова А. З., Турдиев М. Р. БУХОРО ШАҲРИДА МЕЪДА ВА ЁН ИККИ БАРМОҚЛИ ИЧАК ЯРАСИ УЧРАШ ЭПИДЕМИОЛОГИЯСИ //Oriental renaissance: Innovative, educational, natural and social sciences. – 2022. – Т. 2. – №. 4. – С. 642-647.
25. Zokirovna O. A. Modern Concepts of Idiopathic Pulmonary Fibrosis //American Journal of Pediatric Medicine and Health Sciences. – 2023. – Т. 1. – №. 3. – С. 97-101.
26. Zokirovna O. A. Pathology of Precancerous Conditions of the Ovaries //American Journal of Pediatric Medicine and Health Sciences. – 2023. – Т. 1. – №. 3. – С. 93-96.
27. Зокировна, Олимова Азиза и Тешаев Шухрат Джумаевич. «Морфологические аспекты печени белых беспородных крыс после тяжелой черепно-мозговой травмы, вызванной экспериментально в виде дорожно-транспортного происшествия». *Scholastic: Journal of Natural and Medical Education* 2.2 (2023): 59-62.
28. Zokirovna O. A. Comparative characteristics of the morphological parameters of the liver at different periods of traumatic brain injury //Euro-Asia Conferences. – 2021. – С. 139-142.
29. Zokirovna O. A. Macroand microscopic structure of the liver of threemonthly white rats //Academic research in educational sciences. – 2021. – Т. 2. – №. 9. – С. 309-312.
30. Олимова А. З. Частота Встречаемости Миомы Матки У Женщин В Репродуктивном Возрасте //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIY JURNALI. – 2021. – Т. 1. – №. 6. – С. 551-556.
31. Zokirovna O. A., Abdurasulovich S. B. Ovarian Diseases in Age of Reproductive Women: Dermoid Cyst //IJTIMOIY FANLARDA INNOVASIYA ONLAYN ILMIY JURNALI. – 2021. – Т. 1. – №. 6. – С. 154-161.
32. Zokirovna O. A. Cytological screening of cervical diseases: pap test research in the bukhara regional diagnostic center for the period 2015-2019. – 2022.
33. Zokirovna O. A., PREVALENCE R. M. M. EPIDEMIOLOGY OF CANCER OF THE ORAL CAVITY AND THROAT IN THE BUKHARA REGION //Web of Scientist: International Scientific Research Journal. – 2022. – Т. 3. – №. 11. – С. 545-550.

34. Olimova A. Z. The frequency of occurrence of my uterus In women of reproductive age //JOURNAL OF ADVANCED RESEARCH AND STABILITY (JARS). – 2021. – T. 1. – №. 06. – C. 551-556.
35. Olimova Aziza Zokirovna. (2023). MODERN PRINCIPLES OF THE EFFECT OF HEMODIALYSIS THERAPY ON HEART RATE. *International Journal of Integrative and Modern Medicine*, 1(1), 80–85. Retrieved from <http://medicaljournals.eu/index.php/IJIMM/article/view/28>
36. Olimova Aziza Zokirovna. (2023). PATHOMORPHOLOGICAL CHARACTERISTICS OF THE EPIDIDYMIS UNDER IRRADIATION. *International Journal of Integrative and Modern Medicine*, 1(1), 96–100. Retrieved from <http://medicaljournals.eu/index.php/IJIMM/article/view/31>
37. Olimova Aziza Zokirovna. (2023). THE INCIDENCE OF CANCER OF THE ORAL CAVITY AND PHARYNX IN THE BUKHARA REGION. *International Journal of Integrative and Modern Medicine*, 1(1), 86–89. Retrieved from <http://medicaljournals.eu/index.php/IJIMM/article/view/29>
38. Olimova Aziza Zokirovna. (2023). INFLUENCE OF ALCOHOL INTOXICATION ON THE HEART TISSUE OF RATS IN THE EXPERIMENT. *International Journal of Integrative and Modern Medicine*, 1(1), 90–95. Retrieved from <http://medicaljournals.eu/index.php/IJIMM/article/view/30>
39. Olimova Aziza Zokirovna. (2023). Modern Aspects of the Etiology of Gastric Ulcer and Its Complications. *American Journal of Pediatric Medicine and Health Sciences (2993-2149)*, 1(3), 163–166. Retrieved from <http://grnjournal.us/index.php/AJPMHS/article/view/208>
40. Zokirovna O. A., Jumaevich T. S. Morphological Aspects of the Liver of White Outbred Rats After Severe Traumatic Brain Injury Caused Experimentally in the Form of a Road Accident //Scholastic: Journal of Natural and Medical Education. – 2023. – T. 2. – №. 2. – C. 59-62.
41. Aziza Zokirovna Olimova GASTRIC ULCER AND ITS COMPLICATIONS // Scientific progress. 2022. №3. URL: <https://cyberleninka.ru/article/n/gastric-ulcer-and-its-complications> (дата обращения: 28.09.2023).
42. Olimova Aziza Zokirovna. (2022). TECHNIQUE FOR CUTTING BIOPSY AND SURGICAL MATERIAL IN THE PRACTICE OF PATHOLOGICAL ANATOMY AND FORENSIC MEDICINE. *Web of Scientist: International Scientific Research Journal*, 3(7), 116–120. <https://doi.org/10.17605/OSF.IO/PSQ59>
43. Zhumayevich N. F., Zokirovna O. A. PATHOMORPHOLOGY OF GASTRIC CANCER //BARQARORLIK VA YETAKCHI TADQIQOTLAR ONLAYN ILMIIY JURNALI. – 2022. – C. 330-333.
44. Zokirovna O. A. Epidemiological and Etiological Data of Morphogenesis and Pathomorphology of Congenital Heart Diseases in Children //American Journal of Pediatric Medicine and Health Sciences. – 2023. – T. 1. – №. 4. – C. 88-91.